

# Product datasheet for SC333747

## CCL28 (NM\_001301873) Human Untagged Clone

## **Product data:**

#### OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| Product Type:                | Expression Plasmids   |
|------------------------------|---|
| Product Name:                | CCL28 (NM_001301873) Human Untagged Clone   |
| Tag:                         | Tag Free  |
| Symbol:                      | CCL28   |
| Synonyms:                    | CCK1; MEC; SCYA28   |
| Mammalian Cell<br>Selection: | Neomycin  |
| Vector:                      | pCMV6-Entry (PS100001)  |
| E. coli Selection:           | Kanamycin (25 ug/mL)  |
| Fully Sequenced ORF:         | <pre>&gt;SC333747 representing NM_001301873. Blue=Insert sequence Red=Cloning site Green=Tag(s)</pre>   |
|                              | GCTCGTTTAGTGAACCGTCAGAATTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCCGCCGCGCGCGCCCATGCAGCAGAGGAGCTCGCCATCGTGGCCTTGGCTGTCTGT |
| <b>Restriction Sites:</b>    | Sgfl-Mlul   |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### Plasmid Map:



| ACCN:                  | NM_001301873   |
|------------------------|--|
| Insert Size:           | 384 bp   |
| OTI Disclaimer:        | Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).   |
| Components:            | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).   |
| Reconstitution Method: | <ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol> |
| RefSeq:                | <u>NM 001301873.1</u>  |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

| <b>ORIGENE</b> CCL28 (NM_001301873) Human Untagged Clone – SC333747 |   |  |
|---|---|--|
| RefSeq Size:  | 1370 bp   |  |
| RefSeq ORF:   | 384 bp  |  |
| Locus ID:   | 56477   |  |
| UniProt ID:   | <u>Q9NRJ3</u>   |  |
| Cytogenetics:   | 5p12  |  |
| Protein Families:   | Druggable Genome, Secreted Protein  |  |
| Protein Pathways:   | Chemokine signaling pathway, Cytokine-cytokine receptor interaction   |  |
| MW:   | 14.3 kDa  |  |
| Gene Summary:   | This antimicrobial gene belongs to the subfamily of small cytokine CC genes. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for resting CD4 or CD8 T cells and eosinophils. The product of this gene binds to chemokine receptors CCR3 and CCR10. This chemokine may play a role in the physiology of extracutaneous epithelial tissues, including diverse mucosal organs. Multiple transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Sep 2014]<br>Transcript Variant: This variant (2) differs in the 3' UTR compared to variant 1. Variants 1, 2, and 3 all encode the same isoform (a). |  |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US