

Product datasheet for SC333787

PHF19 (NM_001286843) Human Untagged Clone

Product data:

OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| N I I I | |
|------------------------------|---|
| Product Type: | Expression Plasmids |
| Product Name: | PHF19 (NM_001286843) Human Untagged Clone |
| Tag: | Tag Free |
| Symbol: | PHF19 |
| Synonyms: | MTF2L1; PCL3; TDRD19B |
| Mammalian Cell Selection: | Neomycin |
| Vector: | pCMV6-Entry (PS100001) |
| E. coli Selection: | Kanamycin (25 ug/mL) |
| Fully Sequenced ORF: | <pre>>SC333787 representing NM_001286843. Blue=Insert sequence Red=Cloning site Green=Tag(s) GCTCGTTTAGTGAACCGTCAGAATTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTG GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC ATGGAGAATCGAGCTCTGGATCCAGGGACTCGGGACTCCTATGGTGCCACCAGCCACCTCCCCAACAAG GGGGCCCTGGCGAAGGTCAAGAACAACTTCAAAGACTTGATGTCCAAACTGACGGAGGGCCAGTATGTG CTGTGCCGGTGGACAGATGGCCTGTACTACCTCGGGAAGATCAAGAGGGTCAGCAGCACTCAAGAAGC TGCCTCGTGACTTTCGAAGATAATTCCAAATACTGGGTCCTATGGAAGGACATACAGCATGCCGGTGTT</pre> |
| | CCAGGAGAGGAGCCCAAGTGCAACATCTGCCTAGGGAAGACATCAGGGCCGCTGAATGAGATCCTCATC TGCGGGAAGTGTGGCCTGGTGCCACATCCCCATAGCGGGCAGTGCTGA ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC |
| Restriction Sites: | Sgfl-Mlul |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Plasmid Map:



| ACCN: | NM_001286843 |
|------------------------|--|
| Insert Size: | 393 bp |
| OTI Disclaimer: | Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP). |
| Components: | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water). |
| Reconstitution Method: | Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. |
| RefSeq: | <u>NM 001286843.1</u> |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

| | PHF19 (NM_001286843) Human Untagged Clone – SC333787 |
|------------------|--|
| RefSeq Size: | 981 bp |
| RefSeq ORF: | 393 bp |
| Locus ID: | 26147 |
| UniProt ID: | <u>Q5T6S3</u> |
| Cytogenetics: | 9q33.2 |
| Protein Families | : Druggable Genome |
| MW: | 14.4 kDa |
| Gene Summary: | Polycomb group (PcG) that specifically binds histone H3 trimethylated at 'Lys-36' (H3K36me3) and recruits the PRC2 complex. Probably involved in the transition from an active state to a repressed state in embryonic stem cells: acts by binding to H3K36me3, a mark for transcriptional activation, and recruiting H3K36me3 histone demethylases RIOX1 or KDM2B, leading to demethylation of H3K36 and recruitment of the PRC2 complex that mediates H3K27me3 methylation, followed by de novo silencing. Recruits the PRC2 complex to CpG islands and contributes to embryonic stem cell self-renewal. Also binds dimethylated at 'Lys- 36' (H3K36me2). Isoform 1 and isoform 2 inhibit transcription from an HSV-tk promoter. [UniProtKB/Swiss-Prot Function] Transcript Variant: This variant (5) contains alternate exon structure in both the 5' and 3' regions, and it thus differs in both UTRs, initiates translation from a downstream in-frame start codon, and includes an alternate 3' coding region, compared to variant 3. The encoded |

isoform (e) is shorter at the N-terminus and has a distinct C-terminus, compared to isoform c.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US