

# **Product datasheet for SC333809**

## OAZ1 (NM 001301020) Human Untagged Clone

### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** OAZ1 (NM\_001301020) Human Untagged Clone

Tag: Tag Free Symbol: OAZ1

Synonyms: AZ1; AZI; OAZ

Mammalian Cell Selection:

Neomycin

**Vector:** pCMV6-Entry (PS100001) **E. coli Selection:** Kanamycin (25 ug/mL)

Fully Sequenced ORF: >SC333809 representing NM\_001301020.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

GTCTCCCTCCACTGCTGTAGTAACCCGGGTCCGGGGCCTCGGTGGTGCTCC

**ACGCGTACGCGGCCGCTC**GAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT

TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

**Restriction Sites:** Sgfl-Mlul


**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



#### Plasmid Map:



**ACCN:** NM\_001301020

**Insert Size:** 396 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 001301020.1</u>



#### OAZ1 (NM\_001301020) Human Untagged Clone - SC333809

 RefSeq Size:
 1189 bp

 RefSeq ORF:
 396 bp

 Locus ID:
 4946

 Cytogenetics:
 19p13.3

 MW:
 14.3 kDa

**Gene Summary:** The protein encoded by this gene belongs to the ornithine decarboxylase antizyme family,

which plays a role in cell growth and proliferation by regulating intracellular polyamine levels. Expression of antizymes requires +1 ribosomal frameshifting, which is enhanced by high levels of polyamines. Antizymes in turn bind to and inhibit ornithine decarboxylase (ODC), the key enzyme in polyamine biosynthesis; thus, completing the auto-regulatory circuit. This gene encodes antizyme 1, the first member of the antizyme family, that has broad tissue distribution, and negatively regulates intracellular polyamine levels by binding to and targeting ODC for degradation, as well as inhibiting polyamine uptake. Antizyme 1 mRNA contains two potential in-frame AUGs; and studies in rat suggest that alternative use of the two translation initiation sites results in N-terminally distinct protein isoforms with different subcellular localization. Alternatively spliced transcript variants have also been noted for this

Transcript Variant: This variant (2) uses an alternate, in-frame acceptor splice site at the second exon compared to variant 1. The resulting isoform (2) lacks 2 internal amino acids

compared to isoform 1.

gene. [provided by RefSeq, Dec 2014]