

Product datasheet for SC333811

i Todact datasilect for Scoopsoff

UBE2C (NM_001281741) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: UBE2C (NM_001281741) Human Untagged Clone

Tag: Tag Free Symbol: UBE2C

Synonyms: dJ447F3.2; UBCH10

Mammalian Cell

Selection:

Neomycin

Vector: pCMV6-Entry (PS100001) **E. coli Selection:** Kanamycin (25 ug/mL)

Fully Sequenced ORF: >SC333811 representing NM_001281741.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

ATGGCTTCCCAAAACCGCGACCCAGCCGCCACTAGCGTCGCCGCCGCCGCCGTAAAGGAGCTGAGCCGAGCGGGGGGGCGCCGCCGGGGGTCCGGTGGGCAAAAGGCTACAGCAGGAGCTGATGACCCTCATGGCAGTGGGGAGCTCAGAACCAGCTCAACAGTTTGTCTACTGTCCGGTCCCAGAGAAACTCAAGATTCTAGCAAGCCCCTTGTGTGGGGCTTGGGTTGGGACATGAGGCTGCTGCTGGAGCTTACTCTGCAACTGTTTCTCCAAATGCCAGAACCCAACATTGATAGTCCCTTGAACACACTGCTGCCGAGCTCTGGAAAAACCCCACAGCTTTT

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT

TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites: Sgfl-Mlul

ACCN: NM_001281741

Insert Size: 399 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 001281741.1</u>

 RefSeq Size:
 723 bp

 RefSeq ORF:
 399 bp

 Locus ID:
 11065

 UniProt ID:
 000762

 Cytogenetics:
 20q13.12

Protein Families: Druggable Genome

Protein Pathways: Ubiquitin mediated proteolysis

MW: 14.4 kDa

Gene Summary: The modification of proteins with ubiquitin is an important cellular mechanism for targeting

abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, ubiquitin-conjugating enzymes, and ubiquitin-protein ligases. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. The encoded protein is required for the destruction of mitotic cyclins and for cell cycle progression, and may be involved in cancer progression. Multiple transcript variants encoding different isoforms have been found for this gene. Pseudogenes of this gene have been defined on chromosomes 4, 14, 15, 18, and 19. [provided by RefSeq, Aug 2013] Transcript Variant: This variant (7) has multiple differences in the central coding region but

maintains the reading frame, compared to variant 1. The encoded isoform (6) is shorter,

compared to isoform 1.