

## Product datasheet for SC333853

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Host cell factor C1 regulator 1 (HCFC1R1) (NM 001288665) Human Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

Product Name: Host cell factor C1 regulator 1 (HCFC1R1) (NM\_001288665) Human Untagged Clone

Tag: Tag Free

Symbol: HCFC1R1

Synonyms: HPIP

**Vector:** pCMV6-Entry (PS100001)


Fully Sequenced ORF: >SC333853 representing NM\_001288665.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

**TGA** 

**Restriction Sites:** Sgfl-Mlul

Plasmid Map:





ORIGENE

**ACCN:** NM\_001288665

**Insert Size:** 417 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 001288665.1</u>

 RefSeq Size:
 944 bp

 RefSeq ORF:
 417 bp

 Locus ID:
 54985

 UniProt ID:
 Q9NWW0

 Cytogenetics:
 16p13.3

**Protein Families:** Druggable Genome

MW: 15.3 kDa

**Gene Summary:** Regulates HCFC1 activity by modulating its subcellular localization. Overexpression of

HCFC1R1 leads to accumulation of HCFC1 in the cytoplasm. HCFC1R1-mediated export may provide the pool of cytoplasmic HCFC1 required for import of virion-derived VP16 into the

nucleus.[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (4) differs in the 5' UTR, compared to variant 1. Variants 1, 3,

and 4 encode the same isoform (1).