

Product datasheet for SC333892

DIABLO (NM 001278302) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: DIABLO (NM_001278302) Human Untagged Clone

Tag: Tag Free Symbol: DIABLO

Synonyms: DFNA64; SMAC

Mammalian Cell Neomycin

Selection: Vector:

pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

Fully Sequenced ORF: >SC333892 representing NM_001278302.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

CTGCGTGAGGATTGA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGGATCTGGCAGCAAATGATATCCTGGAT

TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites: Sgfl-Mlul




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Plasmid Map:



ACCN: NM_001278302

Insert Size: 429 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001278302.1</u>



DIABLO (NM_001278302) Human Untagged Clone - SC333892

RefSeq Size: 1853 bp
RefSeq ORF: 429 bp
Locus ID: 56616
UniProt ID: Q9NR28
Cytogenetics: 12q24.31

Protein Families: Transmembrane

MW: 16.4 kDa

Gene Summary: This gene encodes an inhibitor of apoptosis protein (IAP)-binding protein. The encoded

mitochondrial protein enters the cytosol when cells undergo apoptosis, and allows activation of caspases by binding to inhibitor of apoptosis proteins. Overexpression of the encoded protein sensitizes tumor cells to apoptosis. A mutation in this gene is associated with young-adult onset of nonsyndromic deafness-64. Alternative splicing results in multiple transcript

variants encoding different isoforms. [provided by RefSeq, May 2013]

Transcript Variant: This variant (5) differs in its 5' UTR and 5' coding region, uses an alternate start codon, and lacks an in-frame exon in the internal coding region, compared to variant 1. The encoded isoform (4) has a distinct N-terminus and is shorter, compared to isoform 1.