

Product datasheet for SC334050

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DNA Primase (PRIM2) (NM_001282488) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: DNA Primase (PRIM2) (NM_001282488) Human Untagged Clone

Tag: Tag Free

Symbol: DNA Primase

Mammalian Cell Neomycin

Selection:

Synonyms:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >SC334050 representing NM_001282488.

p58; PRIM2A

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites: Sgfl-Mlul

ACCN: NM 001282488

Insert Size: 477 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeg: NM 001282488.1

 RefSeq Size:
 902 bp

 RefSeq ORF:
 477 bp

 Locus ID:
 5558

 UniProt ID:
 P49643

 Cytogenetics:
 6p11.2

Protein Pathways: DNA replication, Metabolic pathways, Purine metabolism, Pyrimidine metabolism

MW: 19.1 kDa

Gene Summary: This gene encodes the 58 kilodalton subunit of DNA primase, an enzyme that plays a key role

in the replication of DNA. The encoded protein forms a heterodimer with a 49 kilodalton subunit. This heterodimer functions as a DNA-directed RNA polymerase to synthesize small RNA primers that are used to create Okazaki fragments on the lagging strand of the DNA. Alternative splicing of this gene results in multiple transcript variants. This gene has a related

pseudogene, which is also present on chromosome 6. [provided by RefSeq, Apr 2014] Transcript Variant: This variant (3) differs in the 5' UTR, lacks multiple 3' coding exons, and its 3' terminal exon extends past a splice site used in variant 1, resulting in a distinct 3' coding region and 3' UTR, compared to variant 1. The encoded isoform (b) is shorter and has a distinct C-terminus, compared to isoform a. Variants 2 and 3 encode the same isoform (b).