

Product datasheet for SC334074

TMEM159 (NM_001301771) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: TMEM159 (NM_001301771) Human Untagged Clone

Tag: Tag Free
Symbol: TMEM159

Vector: pCMV6-Entry (PS100001)


Fully Sequenced ORF: >SC334074 representing NM_001301771.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

TGA

Restriction Sites: Sgfl-Mlul

Plasmid Map:



ACCN: NM 001301771



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



TMEM159 (NM_001301771) Human Untagged Clone - SC334074

Insert Size: 486 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001301771.1</u>

 RefSeq Size:
 2041 bp

 RefSeq ORF:
 486 bp

 Locus ID:
 57146

 UniProt ID:
 Q96B96

 Cytogenetics:
 16p12.3

Protein Families: Transmembrane

MW: 17.5 kDa

Gene Summary: Plays an important role in the formation of lipid droplets (LD) which are storage organelles at

the center of lipid and energy homeostasis (PubMed:31708432). In association with BSCL2/seipin, defines the sites of LD formation in the endoplasmic reticulum

(PubMed:31708432).[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (4) lacks an alternate in-frame exon in the 5' coding region,

compared to variant 1. The encoded isoform (2) is shorter than isoform 1. Variants 2, 3, and 4

all encode the same isoform (2).