

Product datasheet for SC334202

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

C22orf25 (TANGO2) (NM_001283235) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: C22orf25 (TANGO2) (NM 001283235) Human Untagged Clone

Tag: Tag Free Symbol: TANGO2

Synonyms: C22orf25; MECRCN

Mammalian Cell

Neomycin

Vector:

E. coli Selection:

Selection:

pCMV6-Entry (PS100001)
Kanamycin (25 ug/mL)

Fully Sequenced ORF:

>NCBI ORF sequence for NM_001283235, the custom clone sequence may differ by one or

more nucleotides

TCTCCCACTGGGAGACCAGAACCTATGAGTTCACACTGCAGAGCTAA

Restriction Sites: Sgfl-Mlul

ACCN: NM 001283235

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 001283235.2</u>, <u>NP 001270164.1</u>

RefSeq Size: 3323 bp
RefSeq ORF: 537 bp
Locus ID: 128989
UniProt ID: Q6ICL3
Cytogenetics: 22q11.21

Gene Summary: This gene belongs to the transport and Golgi organization family, whose members are

predicted to play roles in secretory protein loading in the endoplasmic reticulum. Depletion of this gene in Drosophila S2 cells causes fusion of the Golgi with the ER. In mouse tissue culture cells, this protein co-localizes with a mitochondrially targeted mCherry protein and displays very low levels of co-localization with Golgi and peroxisomes. Allelic variants of this gene are associated with rhabdomyolysis, metabolic crises with encephalopathy, and cardiac

arrhythmia. [provided by RefSeq, Apr 2016]