

Product datasheet for SC334203

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TMED4 (NM_001303059) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: TMED4 (NM_001303059) Human Untagged Clone

Tag: Tag Free
Symbol: TMED4

Synonyms: ERS25; GMP25iso; HNLF; p24a3; p24alpha3

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001303059, the custom clone sequence may differ by one or

more nucleotides

AGATGCGTCACCTCAAGAGCTTCTTTGAGGCCAAGAAGCTGGTG<mark>TAG</mark>

Restriction Sites: Sgfl-Mlul

ACCN: NM 001303059

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001303059.1</u>, <u>NP 001289988.1</u>

RefSeq Size: 2226 bp
RefSeq ORF: 537 bp
Locus ID: 222068
Cytogenetics: 7p13

Protein Families: Transmembrane

Gene Summary: Involved in vesicular protein trafficking, mainly in the early secretory pathway. targeting.

Involved in the maintenance of the Golgi apparatus. Appears to play a role in the biosynthesis of secreted cargo including processing. Involved in endoplasmic reticulum stress response. May play a role in the regulation of heat-shock response and apoptosis (By similarity).

[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (3) lacks an in-frame exon compared to variant 1. The encoded

isoform (3) is shorter than isoform 1.