

## **Product datasheet for SC334205**

## **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

## STAC3 (NM\_001286257) Human Untagged Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** STAC3 (NM\_001286257) Human Untagged Clone

Tag: Tag Free Symbol: STAC3

**Synonyms:** MYPBB; NAM

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM\_001286257, the custom clone sequence may differ by one or

more nucleotides

GCCGCAAGGTGGGGCTGTTTCCCACCGACTTTCTAGAGGAAATTTAG

Restriction Sites: Sgfl-Mlul

**ACCN:** NM 001286257

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 001286257.1, NP 001273186.1</u>

RefSeq Size: 1235 bp
RefSeq ORF: 537 bp
Locus ID: 246329
UniProt ID: Q96MF2
Cytogenetics: 12q13.3

**Gene Summary:** The protein encoded by this gene is a component of the excitation-contraction coupling

machinery of muscles. This protein is a member of the Stac gene family and contains an N-terminal cysteine-rich domain and two SH3 domains. Mutations in this gene are a cause of

Native American myopathy. [provided by RefSeq, Nov 2013]

Transcript Variant: This variant (3) differs in the 5' UTR, lacks a portion of the 5' coding region, and initiates translation at a downstream in-frame start codon, compared to variant 1. The

encoded isoform (c) has a shorter N-terminus than isoform a.