

Product datasheet for SC334244

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Granzyme H (GZMH) (NM_001270780) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Granzyme H (GZMH) (NM 001270780) Human Untagged Clone

Tag: Tag Free

Symbol: Granzyme H

Synonyms: CCP-X; CGL-2; CSP-C; CTLA1; CTSGL2

Mammalian Cell

Selection:

Neomycin

Vector: pCMV6-Entry (PS100001) **E. coli Selection:** Kanamycin (25 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001270780, the custom clone sequence may differ by one or

more nucleotides

Restriction Sites: Sgfl-Mlul

ACCN: NM 001270780

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 001270780.1</u>, <u>NP 001257709.1</u>

RefSeq Size: 795 bp
RefSeq ORF: 549 bp
Locus ID: 2999
Cytogenetics: 14q12

Protein Families: Druggable Genome, Protease

Gene Summary: This gene encodes a member of the peptidase S1 family of serine proteases. Alternative

splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed to generate a chymotrypsin-like protease. This protein is reported to be constitutively expressed in the NK (natural killer) cells of the immune system and may play a role in the cytotoxic arm of the innate immune response by inducing target cell death and by directly cleaving substrates in pathogen-infected cells. This gene is present in a gene cluster with another member of the granzyme subfamily on chromosome 14.

[provided by RefSeq, Nov 2015]

Transcript Variant: This variant (2) uses an alternate in-frame splice site in the coding region compared to variant 1. It encodes isoform 2 which is shorter than isoform 1. This isoform (2) may undergo proteolytic processing similar to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.