

Product datasheet for SC334319

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

COA8 (NM_001302653) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: COA8 (NM 001302653) Human Untagged Clone

Tag: Tag Free Symbol: COA8

Synonyms: APOP; APOP1; APOPT1; C14orf153; MC4DN17

Mammalian Cell

Selection:

Neomycin

Vector: pCMV6-Entry (PS100001) **E. coli Selection:** Kanamycin (25 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001302653, the custom clone sequence may differ by one or

more nucleotides

Restriction Sites: Sgfl-Mlul

ACCN: NM 001302653

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 001302653.1</u>, <u>NP 001289582.1</u>

 RefSeq Size:
 1337 bp

 RefSeq ORF:
 573 bp

 Locus ID:
 84334

 UniProt ID:
 Q96IL0

 Cytogenetics:
 14q32.33

Protein Families: Secreted Protein

Gene Summary: This gene encodes a protein that localizes to the mitochondria, where it stimulates the

release of cytochrome c, thereby promoting programmed cell death. Mutations in this gene have been found in individuals with mitochondrial complex IV deficiency. Alternative splicing

results in multiple transcript variants. [provided by RefSeq, Nov 2014]

Transcript Variant: This variant (3) has a shorter 3' UTR, and contains an alternate exon in the 3' coding region, which results in a frameshift, compared to variant 1. The encoded isoform (3) has a distinct, shorter C-terminus than isoform 1. Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were

based on alignments.