

Product datasheet for SC334339

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RHOH (NM_001278367) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: RHOH (NM_001278367) Human Untagged Clone

Tag: Tag Free Symbol: RHOH

Synonyms: ARHH; TTF
Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001278367, the custom clone sequence may differ by one or

more nucleotides

Restriction Sites: Sgfl-Mlul

ACCN: NM_001278367

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 001278367.1</u>, <u>NP 001265296.1</u>

RefSeq Size:1865 bpRefSeq ORF:576 bpLocus ID:399

UniProt ID: Q15669
Cytogenetics: 4p14

Protein Families: Transcription Factors

Protein Pathways: Leukocyte transendothelial migration

Gene Summary: The protein encoded by this gene is a member of the Ras superfamily of guanosine

triphosphate (GTP)-metabolizing enzymes. The encoded protein is expressed in

hematopoietic cells, where it functions as a negative regulator of cell growth and survival. This gene may be hypermutated or misexpressed in leukemias and lymphomas. Chromosomal

translocations in non-Hodgkin's lymphoma occur between this locus and B-cell

CLL/lymphoma 6 (BCL6) on chromosome 3, leading to the production of fusion transcripts. Alternative splicing in the 5' untranslated region results in multiple transcript variants that

encode the same protein. [provided by RefSeq, May 2013]

Transcript Variant: This variant (10) contains an alternate 5' exon and lacks two exons in the 5'

UTR, compared to variant 1. Variants 1 through 12 encode the same protein.