

## **Product datasheet for SC334381**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

## STOM (NM\_001270526) Human Untagged Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** STOM (NM\_001270526) Human Untagged Clone

Tag: Tag Free Symbol: STOM

**Synonyms:** BND7; EPB7; EPB72

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM\_001270526, the custom clone sequence may differ by one or

more nucleotides

Restriction Sites: Sgfl-Mlul

**ACCN:** NM 001270526

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



AAAGGTGGAGCGTGTGGAAATTAAGGATAAA<mark>TGA</mark>



## **Reconstitution Method:**

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** <u>NM 001270526.1</u>, <u>NP 001257455.1</u>

RefSeq Size: 3107 bp
RefSeq ORF: 594 bp
Locus ID: 2040
Cytogenetics: 9q33.2

**Protein Families:** Transmembrane

**Gene Summary:** This gene encodes a member of a highly conserved family of integral membrane proteins.

The encoded protein localizes to the cell membrane of red blood cells and other cell types, where it may regulate ion channels and transporters. Loss of localization of the encoded protein is associated with hereditary stomatocytosis, a form of hemolytic anemia. There is a pseudogene for this gene on chromosome 6. Alternative splicing results in multiple transcript

variants. [provided by RefSeq, Jul 2012]

Transcript Variant: This variant (3) uses two alternate splice sites in the 3' coding region, which result in a frameshift, compared to variant 1. The encoded isoform (c) has a shorter and distinct C-terminus, compared to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were

based on transcript alignments.