

Product datasheet for **SC334407**

MS4A2 (NM_001256916) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	MS4A2 (NM_001256916) Human Untagged Clone
Tag:	Tag Free
Symbol:	MS4A2
Synonyms:	APY; ATOPY; FCER1B; FCERI; IGEL; IGER; IGHHER; MS4A1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>NCBI ORF sequence for NM_001256916, the custom clone sequence may differ by one or more nucleotides

```
ATGGACACAGAAAGTAATAGGAGAGCAAATCTTGCTCTCCACAGGAGCCTTCCAGTGTGCCTGCATTTG
AAGTCTTGAAATATCTCCCAGGAAGTATCTTCAGGCAGACTATTGAAGTCGGCCTCATCCCCACCACT
GCATACATGGCTGACAGTTTTGAAAAAAGAGCAGGAGTTCCTGGGTTTTCTATTTCTGGAATGTTGTCA
ATTATATCTGAAAGGAGAAATGCAACATATCTGGTGAGAGGAAGCCTGGGAGCAAACACTGCCAGCAGCA
TAGCTGGGGAACGGGAATTACCATCCTGATCATCAACCTGAAGAAGAGCTTGGCCTATATCCACATCCA
CAGTTGCCAGAAATTTTTGAGACCAAGTGCTTTATGGCTTCCTTTTCCACTGAAATTGTAGTGATGATG
CTGTTTCTCACCATTCTGGGACTTGGTAGTGCTGTGCTCACTCACAATCTGTGGAGCTGGGAAGAAGTCA
AAGGAAACAAGGTTCCAGAGGATCGTGTTTATGAAGAATTAACATATATTCAGCTACTTACAGTGAGTT
GGAAGACCCAGGGGAAATGTCTCCTCCATTGATTTAA
```

Restriction Sites:	Sgfl-Mlul
ACCN:	NM_001256916
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001256916.1](#), [NP_001243845.1](#)

RefSeq Size: 3529 bp

RefSeq ORF: 600 bp

Locus ID: 2206

Cytogenetics: 11q12.1

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Asthma, Fc epsilon RI signaling pathway

Gene Summary: The allergic response involves the binding of allergen to receptor-bound IgE followed by cell activation and the release of mediators responsible for the manifestations of allergy. The IgE-receptor, a tetramer composed of an alpha, beta, and 2 disulfide-linked gamma chains, is found on the surface of mast cells and basophils. This gene encodes the beta subunit of the high affinity IgE receptor which is a member of the membrane-spanning 4A gene family. Members of this nascent protein family are characterized by common structural features and similar intron/exon splice boundaries and display unique expression patterns among hematopoietic cells and nonlymphoid tissues. This family member is localized to 11q12, among a cluster of membrane-spanning 4A gene family members. Alternative splicing results in multiple transcript variants encoding distinct proteins. Additional transcript variants have been described but require experimental validation. [provided by RefSeq, Mar 2012]
Transcript Variant: This variant (3) lacks an in-frame exon in the coding region, compared to variant 1, which results in a shorter isoform (3), compared to isoform 1, that traffics to the nuclear membrane. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.