

Product datasheet for SC334435

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DAP1 (DAP) (NM_001291963) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: DAP1 (DAP) (NM_001291963) Human Untagged Clone

Tag: Tag Free Symbol: DAP

Mammalian Cell Neomycin

Selection:

Vector:

pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001291963, the custom clone sequence may differ by one or

more nucleotides

Restriction Sites: Sgfl-Mlul

ACCN: NM_001291963

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

GCAGTCACCTGACGTCCTTGTCTATGGTCTTAAAAACAAGAAGGCACACATTTGA

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: NM 001291963.1, NP 001278892.1

RefSeq Size: 2342 bp
RefSeq ORF: 615 bp
Locus ID: 1611
UniProt ID: P51397
Cytogenetics: 5p15.2

Gene Summary: This gene encodes a basic, proline-rich, 15-kD protein. The protein acts as a positive mediator

of programmed cell death that is induced by interferon-gamma. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by

RefSeq, May 2014]

Transcript Variant: This variant (1) encodes the longer isoform (1).