

Product datasheet for SC334489

riodact datasiicct for Sessarios

ST2 (IL1RL1) (NM 001282408) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: ST2 (IL1RL1) (NM_001282408) Human Untagged Clone

Tag: Tag Free Symbol: IL1RL1

Synonyms: DER4; FIT-1; IL33R; ST2; ST2L; ST2V; T1

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001282408, the custom clone sequence may differ by one or

more nucleotides

TTCTGA

Restriction Sites: Sgfl-Mlul

ACCN: NM_001282408

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: NM 001282408.1, NP 001269337.1

 RefSeq Size:
 3589 bp

 RefSeq ORF:
 636 bp

 Locus ID:
 9173

 UniProt ID:
 Q01638

 Cytogenetics:
 2q12.1

Protein Families: Druggable Genome, Secreted Protein, Transmembrane

Gene Summary: The protein encoded by this gene is a member of the interleukin 1 receptor family. Studies of

the similar gene in mouse suggested that this receptor can be induced by proinflammatory stimuli, and may be involved in the function of helper T cells. This gene, interleukin 1 receptor, type I (IL1R1), interleukin 1 receptor, type II (IL1R2) and interleukin 1 receptor-like 2 (IL1RL2) form a cytokine receptor gene cluster in a region mapped to chromosome 2q12. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Jul

2008]

Transcript Variant: This variant (3) has multiple differences, compared to variant 1. These differences result in distinct 5' and 3' UTRs, cause translation initiation at a downstream start codon and translation termination at a different stop codon, compared to variant 1. It encodes isoform 3 which has a shorter N-terminus and a shorter and distinct C-terminus,

compared to isoform 1.