

## **Product datasheet for SC334527**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

## NBL1 (NM\_001204086) Human Untagged Clone

**Product data:** 

**Product Type:** Expression Plasmids

Product Name: NBL1 (NM\_001204086) Human Untagged Clone

Tag: Tag Free Symbol: NBL1

Synonyms: D1S1733E; DAN; DAND1; NB; NO3

Mammalian Cell Neomycin

Selection:

E. coli Selection:

**Vector:** pCMV6-Entry (PS100001)

Fully Sequenced ORF: >NCBI ORF sequence for NM\_001204086, the custom clone sequence may differ by one or

more nucleotides

Kanamycin (25 ug/mL)

GAGGGGCTGAGGACTGA

**Restriction Sites:** Sgfl-Mlul

**ACCN:** NM\_001204086

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





## **Reconstitution Method:**

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 001204086.2</u>, <u>NP 001191015.1</u>

RefSeq Size: 2178 bp
RefSeq ORF: 648 bp
Locus ID: 4681
Cytogenetics: 1p36.13

**Protein Families:** Secreted Protein

Gene Summary: This gene product is the founding member of the evolutionarily conserved CAN (Cerberus

and DAN) family of proteins, which contain a domain resembling the CTCK (C-terminal cystine knot-like) motif found in a number of signaling molecules. These proteins are secreted, and act as BMP (bone morphogenetic protein) antagonists by binding to BMPs and preventing them from interacting with their receptors. They may thus play an important role during growth and development. Alternatively spliced transcript variants have been identified for this gene. Read-through transcripts between this locus and the upstream mitochondrial inner membrane organizing system 1 gene (GeneID 440574) have been observed. [provided by

RefSeq, May 2013]

Transcript Variant: This variant (5) differs in the 5' UTR, lacks a portion of the 5' coding region, and initiates translation at an alternate start codon, compared to variant 1. The encoded isoform (3) has a distinct N-terminus and is shorter than isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.