

Product datasheet for SC334546

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

FUSIP1 (SRSF10) (NM_001300937) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: FUSIP1 (SRSF10) (NM_001300937) Human Untagged Clone

Tag: Tag Free Symbol: SRSF10

Synonyms: FUSIP1; FUSIP2; NSSR; PPP1R149; SFRS13; SFRS13A; SRp38; SRrp40; TASR; TASR1; TASR2

Mammalian Cell

Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001300937, the custom clone sequence may differ by one or

more nucleotides

Restriction Sites: Sgfl-Mlul

ACCN: NM 001300937

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 001300937.1</u>, <u>NP 001287866.1</u>

RefSeq Size: 4178 bp
RefSeq ORF: 654 bp
Locus ID: 10772
Cytogenetics: 1p36.11

Protein Families: Transcription Factors

Protein Pathways: Spliceosome

Gene Summary: This gene product is a member of the serine-arginine (SR) family of proteins, which are

involved in constitutive and regulated RNA splicing. Members of this family are characterized by N-terminal RNP1 and RNP2 motifs, which are required for binding to RNA, and multiple C-terminal SR/RS repeats, which are important in mediating association with other cellular proteins. This protein interacts with the oncoprotein TLS, and abrogates the influence of TLS on adenovirus E1A pre-mRNA splicing. This gene has pseudogenes on chromosomes 4, 9, 14, 18, and 20. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul

2014]

Transcript Variant: This variant (9) lacks an alternate segment in the 3' coding region, which results in a frameshift, compared to variant 2. The encoded isoform (9) has a distinct C-terminus and is shorter than isoform 2. Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on

alignments.