

Product datasheet for SC334605

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

USP46 (NM_001286768) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: USP46 (NM 001286768) Human Untagged Clone

Tag: Tag Free Symbol: USP46

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001286768, the custom clone sequence may differ by one or

more nucleotides

Restriction Sites: Sgfl-Mlul

ACCN: NM_001286768

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

TTTGATGATGACATTGTAGAGGTTGGTTTGCAGATTATTCTCCAGTAG

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001286768.1</u>, <u>NP 001273697.1</u>

 RefSeq Size:
 8133 bp

 RefSeq ORF:
 678 bp

 Locus ID:
 64854

 UniProt ID:
 P62068

 Cytogenetics:
 4q12

Protein Families: Druggable Genome, Protease

Gene Summary: Modification of cellular proteins by ubiquitin is an essential regulatory mechanism controlled

by the coordinated action of multiple ubiquitin-conjugating and deubiquitinating enzymes. USP46 belongs to a large family of cysteine proteases that function as deubiquitinating enzymes (Quesada et al., 2004 [PubMed 14715245]).[supplied by OMIM, Jun 2009] Transcript Variant: This variant (4) differs in its 5' UTR and initiates translation at a

downstream start codon, compared to variant 1. The encoded isoform (4) has a shorter N-terminus, compared to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on

transcript alignments.