

## **Product datasheet for SC334790**

## Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

EU: info-de@origene.com
CN: techsupport@origene.cn

## PRKACB (NM\_001300917) Human Untagged Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** PRKACB (NM\_001300917) Human Untagged Clone

Tag: Tag Free
Symbol: PRKACB

Synonyms: CAFD2; PKA C-beta; PKACB

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM\_001300917, the custom clone sequence may differ by one or

more nucleotides

**Restriction Sites:** Sgfl-Mlul

**ACCN:** NM 001300917

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).



## PRKACB (NM\_001300917) Human Untagged Clone - SC334790

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001300917.1</u>, <u>NP 001287846.1</u>

 RefSeq Size:
 1961 bp

 RefSeq ORF:
 738 bp

 Locus ID:
 5567

 UniProt ID:
 P22694

 Cytogenetics:
 1p31.1

**Protein Families:** Druggable Genome, Protein Kinase

**Protein Pathways:** Apoptosis, Calcium signaling pathway, Chemokine signaling pathway, Dilated

cardiomyopathy, Gap junction, GnRH signaling pathway, Hedgehog signaling pathway, Insulin

signaling pathway, Long-term potentiation, MAPK signaling pathway, Melanogenesis, Olfactory transduction, Oocyte meiosis, Prion diseases, Progesterone-mediated oocyte maturation, Taste transduction, Vascular smooth muscle contraction, Vibrio cholerae

infection, Wnt signaling pathway

**Gene Summary:** The protein encoded by this gene is a member of the serine/threonine protein kinase family.

The encoded protein is a catalytic subunit of cAMP (cyclic AMP)-dependent protein kinase, which mediates signalling though cAMP. cAMP signaling is important to a number of processes, including cell proliferation and differentiation. Multiple alternatively spliced transcript variants encoding distinct isoforms have been observed. [provided by RefSeq, Jul

2014]

Transcript Variant: This variant (12) differs in both the 5' and 3' exon structures, compared to variant 1. The encoded isoform (12) has distinct N- and C-termini and is shorter than isoform

1.