

## Product datasheet for SC334879

## MED6 (NM 001284211) Human Untagged Clone

**Product data:** 

**Product Type: Expression Plasmids** 

**Product Name:** MED6 (NM\_001284211) Human Untagged Clone

Tag: Tag Free MED6 Symbol:

Synonyms: ARC33; NY-REN-28

**Mammalian Cell** 

Selection:

Neomycin

Vector: pCMV6-Entry (PS100001) E. coli Selection: Kanamycin (25 ug/mL)

>NCBI ORF sequence for NM\_001284211, the custom clone sequence may differ by one or **Fully Sequenced ORF:** 

more nucleotides

ATGGCGGCGGTGGATATCCGAGACAATCTGCTGGGAATTTCTTGGGTTGACAGCTCTTGGATCCCTATTT TGAACAGTGGTAGTGTCCTGGATTACTTTTCAGAAAGAAGTAATCCTTTTTATGACAGAACATGTAATAA TGAAGTGGTCAAAATGCAGAGGCTAACATTAGAACACTTGAATCAGATGGTTGGAATCGAGTACATCCTT TTGCATGCTCAAGAGCCCATTCTTTTCATCATTCGGAAGCAACAGCGGCAGTCCCCTGCCCAAGTTATCC CACTAGCTGATTACTATATCATTGCTGGAGTGATCTATCAGGCACCAGACTTGGGATCAGTTATAAACTC TAGAGTGCTTACTGCAGTGCATGGTATTCAGTCAGCTTTTGATGAAGCTATGTCATACTGTCGATATCAT CCTTCCAAAGGGTATTGGTGGCACTTCAAAGATCATGAAGAGCAAGATAAAGTCAGACCTAAAGCCAAAA GGAAAGAACCAAGCTCTATTTTTCAGAGACAACGTGTGGATGCTTTACTTTTAGACCTCAGACAAAA GTGCTAAAGGCCCCCTGAAAAACGGATGAGACTTCATTACATAATCCTCAGTATTAAAATAAACAGTTG

**CTGA** 

**Restriction Sites:** Sgfl-Mlul

ACCN: NM 001284211

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

> point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## MED6 (NM\_001284211) Human Untagged Clone - SC334879

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001284211.1, NP 001271140.1</u>

 RefSeq Size:
 1700 bp

 RefSeq ORF:
 774 bp

 Locus ID:
 10001

 UniProt ID:
 075586

 Cytogenetics:
 14q24.2

**Protein Families:** Druggable Genome, Transcription Factors

**Gene Summary:** Component of the Mediator complex, a coactivator involved in the regulated transcription of

nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. [UniProtKB/Swiss-Prot

Function]

Transcript Variant: This variant (4) uses an alternate splice site and differs in the 3' exon structure, which results in a frameshift, compared to variant 1. The encoded isoform (4) is

longer and has a distinct C-terminus, compared to isoform 1.