

Product datasheet for SC334895

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Chimaerin 2 (CHN2) (NM_001293077) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Chimaerin 2 (CHN2) (NM 001293077) Human Untagged Clone

Tag: Tag Free

Symbol: Chimaerin 2

Synonyms: ARHGAP3; BCH; CHN2-3; RHOGAP3

Mammalian Cell

Restriction Sites:

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001293077, the custom clone sequence may differ by one or

more nucleotides

ACCN: NM 001293077

Sgfl-Mlul

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).





Chimaerin 2 (CHN2) (NM_001293077) Human Untagged Clone - SC334895

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 001293077.1, NP 001280006.1

 RefSeq Size:
 2799 bp

 RefSeq ORF:
 783 bp

 Locus ID:
 1124

 UniProt ID:
 P52757

 Cytogenetics:
 7p14.3

Gene Summary: This gene encodes a guanosine triphosphate (GTP)-metabolizing protein that contains a

phorbol-ester/diacylglycerol (DAG)-type zinc finger, a Rho-GAP domain, and an SH2 domain. The encoded protein translocates from the cytosol to the Golgi apparatus membrane upon binding by diacylglycerol (DAG). Activity of this protein is important in cell proliferation and migration, and expression changes in this gene have been detected in cancers. A mutation in this gene has also been associated with schizophrenia in men. Alternative transcript splicing and the use of alternative promoters results in multiple transcript variants. [provided by

RefSeg, May 2014]

Transcript Variant: This variant (10, also known as B1-CHNdel ex10-11) lacks two alternate inframe exons, compared to variant 1. The encoded isoform (10) is shorter than isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic

coordinates used for the transcript record were based on transcript alignments.