

## **Product datasheet for SC334989**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

## KIR3DS1 (NM\_001282171) Human Untagged Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** KIR3DS1 (NM\_001282171) Human Untagged Clone

Tag: Tag Free Symbol: KIR3DS1

Synonyms: CD158E2; KIR-123FM; KIR-G1; NKAT-10; NKAT10

**Mammalian Cell** 

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM\_001282171, the custom clone sequence may differ by one or

more nucleotides

CTCCAACAAAAAAAATGCTGCTGTAATGGACCAAGAGCCTGCAGGGAACAGAAG<mark>TGA</mark>

Restriction Sites: Sgfl-Mlul

**ACCN:** NM 001282171

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).





The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube Components:

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

NM 001282171.1, NP 001269100.1 RefSeq:

RefSeq Size: 1446 bp RefSeq ORF: 828 bp Locus ID: 3813 **UniProt ID:** Q14943 Cytogenetics: 19q13.4

**Protein Families:** Transmembrane

**Gene Summary:** Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins expressed

> by natural killer cells and subsets of T cells. The KIR genes are polymorphic and highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among haplotypes, although several "framework" genes are found in all haplotypes (KIR3DL3, KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S) cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for several KIR proteins are subsets of HLA class I molecules; thus, KIR proteins are thought to play an important role in regulation of the immune response. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Aug

Transcript Variant: This variant (3) lacks two consecutive exons in the 5' coding regjon, compared to variant 1. The resulting isoform (3) lacks an internal segment, compared to

isoform 1.

20131