

Product datasheet for SC335035

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

LAIR1 (NM_001289026) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: LAIR1 (NM 001289026) Human Untagged Clone

Tag: Tag Free Symbol: LAIR1

Synonyms: CD305; LAIR-1

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001289026, the custom clone sequence may differ by one or

more nucleotides

TGA

Restriction Sites: Sgfl-Mlul

ACCN: NM 001289026

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).



LAIR1 (NM_001289026) Human Untagged Clone - SC335035

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001289026.2</u>, <u>NP 001275955.2</u>

RefSeq Size:2766 bpRefSeq ORF:843 bpLocus ID:3903

Cytogenetics: 19q13.42

Protein Families: Transmembrane

Gene Summary: The protein encoded by this gene is an inhibitory receptor found on peripheral mononuclear

cells, including natural killer cells, T cells, and B cells. Inhibitory receptors regulate the immune response to prevent lysis of cells recognized as self. The gene is a member of both the immunoglobulin superfamily and the leukocyte-associated inhibitory receptor family. The gene maps to a region of 19q13.4 called the leukocyte receptor cluster, which contains at least 29 genes encoding leukocyte-expressed receptors of the immunoglobulin superfamily. The encoded protein has been identified as an anchor for tyrosine phosphatase SHP-1, and may induce cell death in myeloid leukemias. Alternative splicing results in multiple transcript

variants. [provided by RefSeq, Jan 2014]

Transcript Variant: This variant (f) differs in the 5' UTR, lacks part of the 5' coding region, and uses an alternate start codon, compared to variant a. The encoded isoform (f) has a shorter and distinct N-terminus, compared to isoform a. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record

were based on transcript alignments.