

Product datasheet for SC335053

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RNPS1 (NM_001286626) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: RNPS1 (NM_001286626) Human Untagged Clone

Tag: Tag Free
Symbol: RNPS1
Synonyms: E5.1

Mammalian Cell Neon

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001286626, the custom clone sequence may differ by one or

more nucleotides

TCCCGATAA

Restriction Sites: Sgfl-Mlul

ACCN: NM 001286626

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).



RNPS1 (NM_001286626) Human Untagged Clone - SC335053

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001286626.1</u>, <u>NP 001273555.1</u>

 RefSeq Size:
 2063 bp

 RefSeq ORF:
 849 bp

 Locus ID:
 10921

 UniProt ID:
 Q15287

 Cytogenetics:
 16p13.3

Protein Families: Transcription Factors

Gene Summary: This gene encodes a protein that is part of a post-splicing multiprotein complex involved in

both mRNA nuclear export and mRNA surveillance. mRNA surveillance detects exported mRNAs with truncated open reading frames and initiates nonsense-mediated mRNA decay (NMD). When translation ends upstream from the last exon-exon junction, this triggers NMD to degrade mRNAs containing premature stop codons. This protein binds to the mRNA and remains bound after nuclear export, acting as a nucleocytoplasmic shuttling protein. This protein contains many serine residues. Several transcript variants encoding different isoforms

have been found for this gene. [provided by RefSeq, Nov 2013]

Transcript Variant: This variant (4) lacks an alternate exon compared to variant 2. The resulting

isoform (b) has a shorter and distinct N-terminus compared to isoform a.