

Product datasheet for SC335108

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cyclin Y (CCNY) (NM_001282853) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Cyclin Y (CCNY) (NM_001282853) Human Untagged Clone

Tag: Tag Free Symbol: CCNY

Synonyms: C10orf9; CBCP1; CCNX; CFP1

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001282853, the custom clone sequence may differ by one or

more nucleotides

TGGTCCCCAGCCATCATCTCTTAA

Restriction Sites: Sgfl-Mlul

ACCN: NM_001282853

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).





Cyclin Y (CCNY) (NM_001282853) Human Untagged Clone - SC335108

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001282853.1</u>, <u>NP 001269782.1</u>

 RefSeq Size:
 4774 bp

 RefSeq ORF:
 864 bp

 Locus ID:
 219771

 UniProt ID:
 Q8ND76

 Cytogenetics:
 10p11.21

Gene Summary: Cyclins, such as CCNY, control cell division cycles and regulate cyclin-dependent kinases (e.g.,

CDC2; MIM 116940) (Li et al., 2009 [PubMed 18060517]).[supplied by OMIM, May 2009] Transcript Variant: This variant (4) lacks a portion of the 5' coding region, and initiates

translation at a downstream start codon, compared to variant 1. Variants 2 and 4 encode the same isoform (2), which has a shorter N-terminus than isoform 1. Sequence Note: This RefSeq

record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the

transcript record were based on transcript alignments.