

Product datasheet for SC335113

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

PPP2R2D (NM_001291310) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: PPP2R2D (NM 001291310) Human Untagged Clone

Tag: Tag Free
Symbol: PPP2R2D

Synonyms: B55D; B55delta; MDS026

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001291310, the custom clone sequence may differ by one or

more nucleotides

TACATATTCCAGGACAAAATCAAC<mark>TAG</mark>

Restriction Sites: Sgfl-Mlul

ACCN: NM 001291310

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).





PPP2R2D (NM_001291310) Human Untagged Clone - SC335113

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001291310.1</u>, <u>NP 001278239.1</u>

RefSeq Size: 6791 bp
RefSeq ORF: 867 bp
Locus ID: 55844
UniProt ID: Q66LE6
Cytogenetics: 10q26.3

Protein Families: Druggable Genome, Phosphatase

Protein Pathways: Tight junction

Gene Summary: B regulatory subunit of protein phosphatase 2A (PP2A) that plays a key role in cell cycle by

controlling mitosis entry and exit. The activity of PP2A complexes containing PPP2R2D (PR55-delta) fluctuate during the cell cycle: the activity is high in interphase and low in mitosis. During mitosis, activity of PP2A is inhibited via interaction with phosphorylated ENSA and ARPP19 inhibitors. Within the PP2A complexes, the B regulatory subunits modulate substrate selectivity and catalytic activity, and also may direct the localization of the catalytic enzyme to

a particular subcellular compartment (By similarity).[UniProtKB/Swiss-Prot Function] Transcript Variant: This variant (3) contains an additional exon in the 5' region, and it thus differs in its 5' UTR and initiates translation at a downstream in-frame start codon, compared to variant 1. The encoded isoform (b) is shorter at the N-terminus, compared to isoform a.