

Product datasheet for **SC335144**

ARFGAP1 (NM_001281484) Human Untagged Clone

Product data:

| | |
|---------------------------|--|
| Product Type: | Expression Plasmids |
| Product Name: | ARFGAP1 (NM_001281484) Human Untagged Clone |
| Tag: | Tag Free |
| Symbol: | ARFGAP1 |
| Synonyms: | ARF1GAP; HRIHFB2281 |
| Mammalian Cell Selection: | Neomycin |
| Vector: | pCMV6-Entry (PS100001) |
| E. coli Selection: | Kanamycin (25 ug/mL) |
| Fully Sequenced ORF: | >NCBI ORF sequence for NM_001281484, the custom clone sequence may differ by one or more nucleotides |

```
ATGCTAAGTTCGAGAGTTCCTGGAGTCTCAGGAGGATTACGATCCTTGCTGGTCTTGCAGGAGAAGTA  
CAACAGCAGAGCCGCGGCCCTCTTAGGGATAAGAGTCTCTGGCCAGCCGAGAGTGTGACCGCCTCCTC  
GGACAAGGCTTTTGAAGACTGGCTGAATGATGACCTCGGCTCCTATCAAGGGGCCAGGGGAATCGCTAC  
GTGGGGTTTGGGAACACGCCACCGCCTCAGAAGAAAGAAGATGACTTCTCAACAACGCCATGCTCTCCC  
TGTACTCGGGCTGGAGCAGCTTACCACCTGGAGCCAGCCGGTTTGCCTCGGCAGCCAAGGAGGGCGCTAC  
AAAGTTTGGATCCAAGCGAGTCAGAAGGCGTCCGAGCTGGGCCACAGCCTGAACGAGAACGTCCTCAAG  
CCTGCGCAGGAGAAGGTGAAGGAGGGAAAGATTTTGTATGATGTCTCCAGTGGGGTCTCTCAGTTGGCGT  
CCAAGGTCCAGGGAGTCGGTAGTAAGGGATGGCGGGACGTCACCACCTTTTTTTCGGGAAAGCAGAGGG  
CCCCTTGACAGCCCCTCGGAGGGCCACAGTTATCAGAACAGCGGTCTGGACCACTTCAAAAACAGCAAC  
ATAGACCAGAGCTTCTGGGAGACCTTTGGAAGTCTGAGCCACCAAGACCCGCAAGTCCCCGAGCAGCG  
ACAGCTGGACGTGCGCGGACACCTCCACCAGAGAGGAGGAGCTCGGACAGCTGGGAGGTGTGGGGCTCGGC  
CTCCACCAACAGGAACAGCAACAGCGACGGCGGGGAGGGCGGGGAGGGCACCAGAAGGCAGTGCCGCCG  
GCCGTGCCCACTGATGATGGCTGGGACAACCAGAACTGGTAG
```

| | |
|--------------------|--|
| Restriction Sites: | Sgfl-MluI |
| ACCN: | NM_001281484 |
| OTI Disclaimer: | Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP). |



[View online »](#)

| | |
|-------------------------------|---|
| Components: | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water). |
| Reconstitution Method: | <ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. |
| RefSeq: | <u>NM_001281484.1, NP_001268413.1</u> |
| RefSeq Size: | 3184 bp |
| RefSeq ORF: | 882 bp |
| Locus ID: | 55738 |
| UniProt ID: | <u>Q8N6T3</u> |
| Cytogenetics: | 20q13.33 |
| Protein Pathways: | Endocytosis |
| Gene Summary: | <p>The protein encoded by this gene is a GTPase-activating protein, which associates with the Golgi apparatus and which interacts with ADP-ribosylation factor 1. The encoded protein promotes hydrolysis of ADP-ribosylation factor 1-bound GTP and is required for the dissociation of coat proteins from Golgi-derived membranes and vesicles. Dissociation of the coat proteins is required for the fusion of these vesicles with target compartments. The activity of this protein is stimulated by phosphoinositides and inhibited by phosphatidylcholine. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013]</p> <p>Transcript Variant: This variant (5) lacks two exons in the 5' and central coding region, and initiates translation at an alternate start codon, compared to variant 2. The encoded isoform (e) is shorter and has a distinct N-terminus, compared to isoform b.</p> |