

## **Product datasheet for SC335219**

## PRKACB (NM 001300916) Human Untagged Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** PRKACB (NM\_001300916) Human Untagged Clone

Tag: Tag Free
Symbol: PRKACB

Synonyms: CAFD2; PKA C-beta; PKACB

Mammalian Cell Neomycin

Selection:

Vector:

pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

Fully Sequenced ORF: >SC335219 representing NM\_001300916.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

**ACGCGTACGCGGCCGCTC**GAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT

TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

**Restriction Sites:** Sgfl-Mlul

**ACCN:** NM\_001300916

**Insert Size:** 915 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

## PRKACB (NM\_001300916) Human Untagged Clone - SC335219

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

> point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 001300916.1

RefSeg Size: 2023 bp RefSeq ORF: 915 bp Locus ID: 5567 **UniProt ID:** P22694

Cytogenetics:

1p31.1 **Protein Families:** Druggable Genome, Protein Kinase

**Protein Pathways:** Apoptosis, Calcium signaling pathway, Chemokine signaling pathway, Dilated

cardiomyopathy, Gap junction, GnRH signaling pathway, Hedgehog signaling pathway, Insulin

signaling pathway, Long-term potentiation, MAPK signaling pathway, Melanogenesis, Olfactory transduction, Oocyte meiosis, Prion diseases, Progesterone-mediated oocyte maturation, Taste transduction, Vascular smooth muscle contraction, Vibrio cholerae

infection, Wnt signaling pathway

MW: 35.3 kDa

**Gene Summary:** The protein encoded by this gene is a member of the serine/threonine protein kinase family.

> The encoded protein is a catalytic subunit of cAMP (cyclic AMP)-dependent protein kinase, which mediates signalling though cAMP. cAMP signaling is important to a number of processes, including cell proliferaton and differentiation. Multiple alternatively spliced transcript variants encoding distinct isoforms have been observed. [provided by RefSeq, Jul

2014]

Transcript Variant: This variant (11) differs in the 3' exon structure, compared to variant 1. The

encoded isoform (11) has a distinct C-terminus and is shorter than isoform 1.