

# **Product datasheet for SC335365**

## P2Y6 (P2RY6) (NM 001277206) Human Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** P2Y6 (P2RY6) (NM\_001277206) Human Untagged Clone

Tag:Tag FreeSymbol:P2RY6Synonyms:P2Y6

**Vector:** pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC335365 representing NM\_001277206.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

TGGCAGAGGCAGGGTCGCTGA

Restriction Sites: Sgfl-Mlul




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



### Plasmid Map:



**ACCN:** NM\_001277206

**Insert Size:** 987 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** NM 001277206.1



### P2Y6 (P2RY6) (NM\_001277206) Human Untagged Clone - SC335365

RefSeq Size: 2457 bp
RefSeq ORF: 987 bp
Locus ID: 5031
UniProt ID: Q15077
Cytogenetics: 11q13.4

**Protein Families:** Druggable Genome, GPCR, Transmembrane

**Protein Pathways:** Neuroactive ligand-receptor interaction

MW: 36.4 kDa

**Gene Summary:** The product of this gene belongs to the family of P2 receptors, which is activated by

extracellular nucleotides and subdivided into P2X ligand-gated ion channels and P2Y G-protein coupled receptors. This family has several receptor subtypes with different

pharmacological selectivity, which overlaps in some cases, for various adenosine and uridine nucleotides. This receptor, which is a G-protein coupled receptor, is responsive to UDP, partially responsive to UTP and ADP, and not responsive to ATP. It is proposed that this receptor mediates inflammatory responses. Alternative splicing results in multiple transcript

variants that encode different protein isoforms. [provided by RefSeq, Mar 2013] Transcript Variant: This variant (7) differs in the 5' UTR compared to variant 1. Variants 1,2,3,5,6,7, and 8 encode the same isoform (1). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record

were based on transcript alignments.