

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for SC335375

HSPC142 (BABAM1) (NM_001288756) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	HSPC142 (BABAM1) (NM_001288756) Human Untagged Clone
Tag:	Tag Free
Symbol:	HSPC142
Synonyms:	C19orf62; HSPC142; MERIT40; NBA1
Vector:	pCMV6-Entry (PS100001)
Fully Sequenced ORF:	>SC335375 representing NM_001288756. Blue=Insert sequence <mark>Red</mark> =Cloning site Green=Tag(s)
	ATGGAAGTGGCAGAGCCCAGCAGCCCACTGAAGAGGAGGAGGAGGAAGAGAGGAGCACTCGGCAGAGACCT CGGCCCCGCACTCGCTCCAATCCTGAAGGGGCTGAGGACCGGGCAGTAGGGGCACAGGCCAGCGGCGC AGCCGCAGCGAGGGTGAGGGCGCCGCCAGTGCTGATGATGGGAGCCTCAACACTTCAGGAGCCGGC CCTAAGTCCTGGCAGGTGCCCCCGCCAGCCCCTGAGGTCCAAATTCGGAACCCAAGGGTCAACTGTCCA GAGAAAGTGATTATCTGCCTGGACCTGTCAGAGGAAATGTCACTGCCAAAGCTGGAGTCGTTCAACGGC TCCAAAACCAACGCCCTCAATGTCTCCCAGAAGATGATTGAGATGTTCGTGCGGACAAAACACAAGATC GACAAAAGCCACGAGTTTGCACTGGTGGTGGTGAACGATGACCACGGCCTGGCTGTCTGGCCTGACCTCC GACCACCGCGGAGCTCTGTAGCTGCTCTATGATCTGGAGACGGCCTCCTGTTCCACCTTCAATCTGGAA GGACTTTTCAGCCTCATCCAGCAGAAAACTGAGCTTCCGGTCACAGAGAACGTGCAGACGATTCCCCCG CCATATGTGGTCCGCACCATCCTTGTCTACAGCCGTCCACCTTGCCAGCCCCAGTTCTCCTTGACGGAG CCCATGAAGAAAATGTTCCAGTGGCCCATATTTCTTCTTGACGTTGTTTACATCCACAATGGCACTGAG GAGAAGGAGGAGGAGATGAGTTGGAAGGATATGTTTGCCTTCATGGCAGCCTGGATACCAAGGGTACC AGCTACAAGTATGAGGTGGCACTGGCTGGGCCAGCCCTGGAGTGCACAACTGCAGAGGTACC AGCTACAAGTATGAGGTGGCACTGGCTGGGCCAGCCCTGGAGTGCACAACTGCAAACGGAGGACC AGCTACAAGTATGAGGTGGCACTGGCTGGGCCAGCCCTGGAGTTGCACAACTGCAAACTGGCAAACTGTTG GCCCACCCCCTGCAGCGGCCTTGCCAGAGCCATGCTTCCTACAGCCTGCTGGAGGAGGAGGAGGAGGAGGAGGAGGATGAGCC ATTGAGGTTGAGGCCACTGCTGCAGAGCCATGCTTCCTACAGCCTGCTGGAGGAGGAGGAGGAGGAGGATGAACC ATTGAGGTTGAGGCCACTGCTGCAGACCATGCTTCCTACAGCCTGCTGGAGGAGGAGGAGGAGGATGAACC ATTGAGGTTGAGGCCACTGCTTGCA
Restriction Sites:	Sgfl-Mlul
ACCN:	NM_001288756
Insert Size:	990 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE HSPC142 (BABAM1) (NM_001288756) Human Untagged Clone – SC335375

Reconstitution Method:	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 001288756.1</u>
RefSeq Size:	1456 bp
RefSeq ORF:	990 bp
Locus ID:	29086
UniProt ID:	<u>Q9NWV8</u>
Cytogenetics:	19p13.11
MW:	36.6 kDa
Gene Summary:	Component of the BRCA1-A complex, a complex that specifically recognizes 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1- BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs). The BRCA1-A complex also possesses deubiquitinase activity that specifically removes 'Lys-63'-linked ubiquitin on histones H2A and H2AX. In the BRCA1-A complex, it is required for the complex

integrity and its localization at DSBs. Component of the BRISC complex, a multiprotein

(PubMed:24075985, PubMed:26195665). In these 2 complexes, it is probably required to maintain the stability of BABAM2 and help the 'Lys-63'-linked deubiquitinase activity

mediated by BRCC3/BRCC36 component. The BRISC complex is required for normal mitotic spindle assembly and microtubule attachment to kinetochores via its role in deubiquitinating

deubiquitination of the interferon receptor IFNAR1; deubiquitination increases IFNAR1 activity by enhancing its stability and cell surface expression (PubMed:24075985). Down-regulates the

Transcript Variant: This variant (3) differs in the 5' UTR, compared to variant 1. Variants 1, 2

complex that specifically cleaves 'Lys-63'-linked ubiquitin in various substrates

NUMA1 (PubMed:26195665). Plays a role in interferon signaling via its role in the

(PubMed:24075985).[UniProtKB/Swiss-Prot Function]

and 3 encode the same isoform (1).

response to bacterial lipopolysaccharide (LPS) via its role in IFNAR1 deubiguitination

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US