

Product datasheet for **SC335453**

TXNRD2 (NM_001282512) Human Untagged Clone

Product data:


Product Type:	Expression Plasmids
Product Name:	TXNRD2 (NM_001282512) Human Untagged Clone
Tag:	Tag Free
Symbol:	TXNRD2
Synonyms:	GCCD5; SELZ; TR; TR-BETA; TR3; TRXR2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC335453 representing NM_001282512. Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
GCTCGTTTGTAGTGAACCGTCAGAATTTTGTAAACGACTACTATAGGGCGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGGCGGCAATGGCGGTGGCGCTGCGGGGATTAGGAGGGCGCTTCCGGTGGCGGACGCAGGCCGTGGCG
GGCGGGGTGCGGGGCGCGGCGGGGCGCAGCAGGTCAGCGGGACTATGATCTCCTGGTGGTCGGC
GGGGATCTGGTGGCCTGGCTTGTGCCAAGGAGGCCGCCAGCTGGGAAGGAAGGTGGCCGTGGTGGAC
TACGTGGAACCTTCTCCCAAGGCACCCGGTGGGGCTCGGCGGCACCTGCGTCAACGTGGGCTGCATC
CCCAAGAAGCTGATGCACCAGGCGCAGTCTGGGAGGCCTGATCAAGATGCCCAACTATGGCTGG
GAGGTGGCCAGCCCGTCCGCATGACTGGAGGAAGATGGCAGAAGCTGTTCAAAATCACGTGAAATCC
TTGAACTGGGGCCACCGTGTCCAGCTTCAGGACAGAAAAGTCAAGTACTTTAACATCAAAGCCAGCTTT
GTTGACGAGCACACGGTTTGCGGCGTTGCCAAAGTGGGAAAGAGATTCTGCTGTGAGCCGATCACATC
ATCATTGCTACTGGAGGGCGGCCGAGATACCCACGCACATCGAAGGTGCCTTGGAAATGGAATCACA
AGTGATGACATCTTCTGGCTGAAGGAATCCCTGGAAAAACGTTGGTGGTCGGGGCCAGCTATGTGGCC
CTGGAGTGTGCTGGCTTCTCACCAGGATTGGGCTGGACACCACCATCATGATGCGCAGCATCCCCCTC
CGCGGCTTCGACCAGCAAATGTCCTCCATGGTCATAGAGCACATGGCATCTCATGGCACCCGGTTCCTG
AGGGCTGTGCCCTCGCGGTCAGGAGGCTCCCTGATGGCCAGCTGCAGGTACCTGGGAGGACAGC
ACCACCGCAAGGAGGACACGGGCACCTTTGACACCGTCTGTGGCCATAGCACCTTGATCTCTGGC
TGCTCTCCCAACCACCGTGGGACATGCTGAAAAAACAGAGAAGAGACTGA
ACGCGTACGCGGGCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
```

Restriction Sites: SgfI-MluI



[View online »](#)

Plasmid Map:


ACCN: NM_001282512

Insert Size: 1017 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001282512.1](#)

RefSeq Size: 2081 bp

RefSeq ORF: 1017 bp

Locus ID: 10587

Cytogenetics: 22q11.21

Protein Families: Druggable Genome

Protein Pathways: Pyrimidine metabolism

MW: 36.3 kDa

Gene Summary: The protein encoded by this gene belongs to the pyridine nucleotide-disulfide oxidoreductase family, and is a member of the thioredoxin (Trx) system. Three thioredoxin reductase (TrxR) isozymes are found in mammals. TrxRs are selenocysteine-containing flavoenzymes, which reduce thioredoxins, as well as other substrates, and play a key role in redox homeostasis. This gene encodes a mitochondrial form important for scavenging reactive oxygen species in mitochondria. It functions as a homodimer containing FAD, and selenocysteine (Sec) at the active site. Sec is encoded by UGA codon that normally signals translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, the Sec insertion sequence (SECIS) element, which is necessary for the recognition of UGA as a Sec codon rather than as a stop signal. Alternatively spliced transcript variants encoding different isoforms, including a few localized in the cytosol and some lacking the C-terminal Sec residue, have been found for this gene. [provided by RefSeq, Jun 2017]

Transcript Variant: This variant (5) lacks several exons in the 3' coding region, and contains an alternate 3' terminal exon compared to variant 1. The resulting isoform (5) is shorter, with a distinct C-terminus (lacking selenocysteine) compared to isoform 1.