OriGene Technologies, Inc.
9620 Medical Center Drive, Ste 200
Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Product datasheet for SC335681

PHF19 (NM_001286842) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	PHF19 (NM_001286842) Human Untagged Clone
Tag:	Tag Free
Symbol:	PHF19
Synonyms:	MTF2L1; PCL3; TDRD19B
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC335681 representing NM_001286842. Blue=Insert sequence Red=Cloning site Green=Tag(s) GCTCGTTTAGTGAACCGTCAGAATTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTG GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC ATGCTGCAATGTTACCGGTGCAGGCAGTGGTTCCACGAGGCCTGCACCCAGTGCCTCAATGAGCCCATG ATGTTTGGAGACCGGTTTTACCTGTTCTTCTGCTCCGTGTGTAACCAGGGCCCAGAGTACATCGAGAGG CTGCCCCTGCGATGGGTGGATGTGGTTCACCTGGCCCTCTATAATCTGGGGGTACAGAGCAAGAAGAAG TACTTTGACTTTGAGGAGATTCTGGCCTTTGTCAACCACCACTGGGAGCTCCTGCAGCTTGGCAAGCTC ACCAGCACCCCAGTGACAGATCGAGGACCACATCTCCTCAACGCTCTGAACAGTTATAAAAGCCGGTTC CTCTGCGGCAAGGAGATCAAGAAGAAGAAGTGCATCTTCCGCCTGCGCATCCGCGTCCCACCCAACCCG CCAGGGAAGCTGCTGCCTGACAAAGGACTGCTGCCAAATGAGAACAGCGCCTCCTCTGAGCTGCGTAAG AGAGGAAAGAGCAAGCCTGGTTTGTTGCCTCACGAATTCCAGCAGCAGAAAAGGCGAGTTTATAGAAGA AAAAGATCAAAGTTTTTGCTGGAAGATGCTATTCCCAGTAGTGACTTCACCTCAGCCTGGAGCACCAAC CACCACCTGGCTAGCATATTTGACTTCACGCTGGATGAAATTCAAAGTTTAAAAAGTGCCAGCTCAGGC CAGACCTTCTTCTCAGATGTCGACTCCACCGACGCTGCCAGCACCTCTGGCTCTGCCTCCACCAGCCTC TCCTATGACTCCAGATGGACAGTGGGCAGCCGAAAGAGGAAGCTGGCAGCCAAGGCATACATGCCCCTG CGGGCAAAGCGGTGGGCAGCTGAGCTGGATGGACGCTGCCCCTCGGACAGCAGTGCAGAGGGGGCTTCA GTCCCCGAGCGGCCAGACGAAGGCATTGACAGCCACACATTTGAGAGCATCAGTGAAGATGACTCATCC CTGTCCCACCTCAAGTCATCTATCACCAACTACTTTGGTGCAGCTGGGCGGTTGGCCTGTGGGGAGAAG TACCAGGTGTTGGCTCGGAGGGTCACACCTGAGGGCAAGGTTCAGTACCTGGTGGAGTGGGAAGGGACC ACCCCTTACTGA ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
Restriction Sites:	SgfI-MluI

View online »

Plasmid Map:



ACCN:	NM_001286842
Insert Size:	1116 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001286842.1
RefSeq Size:	3547 bp
RefSeq ORF:	1116 bp
Locus ID:	26147
Cytogenetics:	9q33.2
Protein Families:	Druggable Genome
MW:	42.2 kDa

Gene Summary: Polycomb group (PcG) that specifically binds histone H3 trimethylated at 'Lys-36' (H3K36me3) and recruits the PRC2 complex. Probably involved in the transition from an active state to a repressed state in embryonic stem cells: acts by binding to H3K36me3, a mark for transcriptional activation, and recruiting H3K36me3 histone demethylases RIOX1 or KDM2B, leading to demethylation of H3K36 and recruitment of the PRC2 complex that mediates H3K27me3 methylation, followed by de novo silencing. Recruits the PRC2 complex to CpG islands and contributes to embryonic stem cell self-renewal. Also binds dimethylated at 'Lys-36' (H3K36me2). Isoform 1 and isoform 2 inhibit transcription from an HSV-tk promoter. [UniProtKB/Swiss-Prot Function]
Transcript Variant: This variant (4) lacks several 5' exons but includes an alternate 5' terminal exon, and it thus differs in its 5' UTR and initiates translation from a downstream in-frame start codon, compared to variant 3. The encoded isoform (d) is shorter at the N-terminus, compared to isoform c.