

Product datasheet for **SC335830**

HNF 4 alpha (HNF4A) (NM_001287184) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	HNF 4 alpha (HNF4A) (NM_001287184) Human Untagged Clone
Tag:	Tag Free
Symbol:	HNF4A
Synonyms:	FRTS4; HNF4; HNF4a7; HNF4a8; HNF4a9; HNF4alpha; MODY; MODY1; NR2A1; NR2A21; TCF; TCF-14; TCF14
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC335830 representing NM_001287184. Blue=Insert sequence Red=Cloning site Green=Tag(s)


```
GCTCGTTTAGTGAACCGTCAGAATTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGTCGGACTGGGGCCAGGGCTTCCCCCAGGACCCACCAGACACGTCCCCATCAGAAGGCACCAACCTC
AACCGGCCAACAGCCTGGGTGTGACGCGCCTGTGTGCCATCTGCGGGGACCGGGCCACGGGCAACAC
TACGGTGCCTCGAGCTGTGACGGCTGCAAGGGCTTCTTCCGGAGGAGCGTGCGGAAGAACCACATGTAC
TCCTGCAGATTTAGCCGGCAGTGCCTGGTGGACAAAGACAAGAGGAACCAGTGCCGCTACTGCAGGCTC
AAGAAATGCTTCCGGGCTGGCATGAAGAAGGAAGCCGTCAGAATGAGCGGGACCGGATCAGCACTCGA
AGGTCAAGCTATGAGGACAGCAGCTGCCCTCCATCAATGCGCTCCTGCAGGCGGAGGTCTGTCCCGA
CAGATCACCTCCCCGTCTCCGGATCAACGGCGACATTCGGGCGAAGAAGATTGCCAGCATCGCAGAT
GTGTGTGAGTCCATGAAGGAGCAGCTGCTGTTCTCGTTGAGTGGCCAAGTACATCCCAGCTTTCTGC
GAGTCCCCCTGGACGACAGGTGGCCCTGCTCAGAGCCCATGCTGGCGAGCACCTGCTGCTCGGAGCC
ACCAAGAGATCCATGGTGTTCAGGACGTGCTGCTCCTAGGCAATGACTACATTGCTCCCTCGGCACTGC
CCGGAGCTGGCGGAGATGAGCCGGGTGCCATACGCATCCTTGACGAGCTGGTGTGCCCTTCCAGGAG
CTGCAGATCGATGACAATGAGTATGCCTACCTCAAAGCCATCATCTTCTTTGACCAGATGCCAAGGGG
CTGAGCGATCCAGGAAGATCAAGCGGCTGCGTTCAGGTGCAGGTGAGCTTGGAGGACTACATCAAC
GACCGCCAGTATGACTCGCGTGGCCGCTTTGGAGAGCTGCTGCTGCTGCTGCCACCTTGACAGAGCATC
ACCTGGCAGATGATCGAGCAGATCCAGTTCATCAAGCTCTTCGGCATGGCCAAGATTGACAACTGTTG
CAGGAGATGCTGCTGGGAGTCCGTGCCAAGCCAGGAGGGGCGGGTGGAGTGGGACTCCCCAGGA
GACAGGCCTCACACAGTGAAGTCAACCCCTCAGCTCCTTGCTTCCCCACTGTGCCCTTTGGCAAGTT
GCTTAA
ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
```



[View online »](#)

Restriction Sites: Sgfl-MluI

Plasmid Map:



ACCN: NM_001287184

Insert Size: 1179 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001287184.1](#)

RefSeq Size: 1671 bp

RefSeq ORF: 1179 bp

Locus ID: 3172

UniProt ID: [P41235](#)

Cytogenetics: 20q13.12

Protein Families:	Druggable Genome, ES Cell Differentiation/IPS, Nuclear Hormone Receptor, Transcription Factors
Protein Pathways:	Maturity onset diabetes of the young
MW:	43.8 kDa
Gene Summary:	<p>The protein encoded by this gene is a nuclear transcription factor which binds DNA as a homodimer. The encoded protein controls the expression of several genes, including hepatocyte nuclear factor 1 alpha, a transcription factor which regulates the expression of several hepatic genes. This gene may play a role in development of the liver, kidney, and intestines. Mutations in this gene have been associated with monogenic autosomal dominant non-insulin-dependent diabetes mellitus type I. Alternative splicing of this gene results in multiple transcript variants encoding several different isoforms. [provided by RefSeq, Apr 2012]</p> <p>Transcript Variant: This variant (10) differs at the 5' end (resulting in translation initiation from an alternate upstream start codon) and at the 3' end (that causes a frame-shift) compared to variant 2. The resulting shorter isoform (10, also known as HNF4alpha12) has distinct N- and C-termini compared to isoform 2.</p>