

Product datasheet for **SC335850**

CD36 (NM_001289911) Human Untagged Clone

Product data:


Product Type:	Expression Plasmids
Product Name:	CD36 (NM_001289911) Human Untagged Clone
Tag:	Tag Free
Symbol:	CD36
Synonyms:	BDPLT10; CHDS7; FAT; GP3B; GP4; GPIV; PASIV; SCARB3
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC335850 representing NM_001289911. Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
GCTCGTTTGTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGCCGGGAATTCGTGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGATGAACAGCAGCAACATTCAAGTTAAGCAAAGAGGTCCTTATACGTACAGAGTTCGTTTTCTAGCC
AAGGAAAATGTAACCCAGGACGCTGAGGACAACACAGTCTCTTTCCTGCAGCCCAATGGTGCCATCTTC
GAACCTTCACTATCAGTTGGAACAGAGGCTGACAACTTACAGTTCTCAATCTGGCTGTGGCAGCTGCA
TCCCATATCTATCAAAATCAATTTGTTCAAATGATCCTCAATTCACTTATTAACAAGTCAAAATCTTCT
ATGTTCCAAGTCAGAATTTGAGAGAAGTGTATGGGGCTATAGGGATCCATTTTGTAGTTTGGTCCG
TACCCTGTACTACCACAGTTGGTCTGTTTTATCCTTACAACAATACTGCAGATGGAGTTTATAAAGTT
TTCAATGGAAAAGATAACATAAGTAAAGTTGCCATAATCGACACATATAAAGGTAAGGAATCTGTCC
TATTGGGAAAGTCACTGCGACATGATTAATGGTACAGATGCAGCCTCATTTCACCTTTTGTGAGAAA
AGCCAGGTATTGCAGTCTTTTCTTCTGATATTTGCAGGTCAATCTATGTGATTTGAATCCGACGTT
AATCTGAAAGGAATCCCTGTGTATAGATTTGTTCTTCCATCCAAGGCCTTGCCTCTCCAGTTGAAAAC
CCAGACAATATTGTTTCTGCACAGAAAAAATATCTCAAAAAATGTACATCATATGGTGTGCTAGAC
ATCAGCAAATGCAAAGAAGGGAGACCTGTGTACATTTCACTTCTCATTCTGTATGCAAGTCTGAT
GTTTCAGAACCTATTGATGGATTAACCCAAATGAAGAAGAACATAGGACATACTGGATATTGAACCT
ATAACTGGATTCACTTTACAATTTGCAAAACGGCTGCAGGTCAACCTATTGGTCAAGCCATCAGAAAAA
ATTCAAGTATTAAGAATCTGAAGAGGAAGTATATTGTGCTATTCTTTGGCTTAATGAGACTGGGACC
ATTGGTGTGAGAAAGCAACATGTTTCAGAAAGTCAAGTAACTGGAAAAATAAACCTCCTTGGCCTGATA
GAAATGATCTTACTCAGTGTGGTGTGGTGTGTTTGTGCTTTTATGATTTTATATTGTGCATGCAGA
TCGAAAACAATAAAATAA
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
```

Restriction Sites: Sgfl-Mlul



[View online >](#)

Plasmid Map:


ACCN: NM_001289911

Insert Size: 1191 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001289911.1](#)

RefSeq Size: 1747 bp

RefSeq ORF: 1191 bp

Locus ID: 948

UniProt ID: [P16671](#)

Cytogenetics: 7q21.11

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Adipocytokine signaling pathway, ECM-receptor interaction, Hematopoietic cell lineage, PPAR signaling pathway

MW: 44.7 kDa

Gene Summary: The protein encoded by this gene is the fourth major glycoprotein of the platelet surface and serves as a receptor for thrombospondin in platelets and various cell lines. Since thrombospondins are widely distributed proteins involved in a variety of adhesive processes, this protein may have important functions as a cell adhesion molecule. It binds to collagen, thrombospondin, anionic phospholipids and oxidized LDL. It directly mediates cytoadherence of Plasmodium falciparum parasitized erythrocytes and it binds long chain fatty acids and may function in the transport and/or as a regulator of fatty acid transport. Mutations in this gene cause platelet glycoprotein deficiency. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Feb 2014]

Transcript Variant: This variant (8) contains an alternate 5' UTR, lacks a portion of the 5' coding region, and initiates translation at an alternate downstream AUG, compared to variant 3. The resulting isoform (4) has a shorter N-terminus, compared to isoform 1.