

Product datasheet for **SC335885**

ABI2 (NM_001282927) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	ABI2 (NM_001282927) Human Untagged Clone
Tag:	Tag Free
Symbol:	ABI2
Synonyms:	ABI-2; ABI2B; AbIBP3; AIP-1; AIP1; argBP1; argBP1A; argBPIB; SSH3BP2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC335885 representing NM_001282927. Blue=Insert sequence Red=Cloning site Green=Tag(s)


```
GCTCGTTTAGTGAACCGTCAGAATTTGTAAACGACTACTATAGGGCGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCCGCATCGCC
ATGTCCTGCAGATGCTGGATATCCAGGCATCCCAGCTACGAAGGATGGAATCTTCAATCAATCATATTT
CACAAGCAAATTAGAGGCGTTGATCTTGAGTCGACTTTTGTACCAAATTTGAAACAATTCGAGTTTG
AGATTGAATGAGACAGTTGATATTCATAAAGAGAAAGTTGCAAGAAGAGAAATGGTATTTTGACTACC
AATAAAAAACACTCAAGGACACATAAGATTATTGCTCCAGCCAACCTTGAACGACCAGTTCGTTATATT
AGAAAACCTATTGACTATACAATTCTAGATGATATTGGACATGGAGTAAAGGTGAGTACCCAGAACATG
AAGATGGGTGGGCTGCCCGTACAACACCTCCAACCTCAGAAGCCCCTAGTCCCCTATGTCAGGGAAA
GGGACACTTGGGCGGCACTCCCCCTATCGCACACTGGAGCCAGTGGCTCCTCCAGTGGTACCAAATGAT
TACGTACCTAGCCCAACCCGTAATATGGCTCCCTCGCAGCAGAGCCCTGTGAGGACAGCTTCTGTGAAT
CAAAGAAATCGAACTTACAGCAGCAGTGGGAGTAGTGGAGGGAGCCACCAAGTAGTCGGAGCAGCAGT
CGAGAGAACAGTGAAGTGGTAGTGTGGGGTTCTATTGCTGTTCTACTCCATCTCTCCCAAGTGC
TTTCCAGGTCATCCTGTACAGTTCTACAGCATGAATAGGCCTGCCTCTCGCCATACTCCCCAACATA
GGGGGCTCGTTGCCCTATAGACGCCCTCCTTCCATTACTTCACAAACAAGCCTTCAGAATCAGATGAAT
GGAGGACCTTTTATAGCCAGAATCCAGTTTCAGATACACCACCTCCACCGCCACCTGTGGAAGAACCA
GTCTTTGATGAGTCTCCCCACCTCCTCCTCCAGAAGATTACGAAGAGGAGGAAGCTGTGGTT
GAGTATAGTGATCCTTATGCTGAAGAGGACCCACCGTGGGCTCCAGTTCTTACTTGAAAAGGTTGTG
GCAATTTATGACTATACAAAAGACAAGGAAGATGAGCTGTCTTTTCAGGAAGGAGCCATTATTTATGTC
ATCAAGAAGAATGACGATGGTTGGTATGAGGGAGTTATGAATGGAGTGACTGGGCTTTTTCTGGGAAT
TACGTTGAGTCTATCATGCATTATTCTGAGTAA
ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
```

Restriction Sites: Sgfl-MluI



[View online »](#)

Plasmid Map:



ACCN: NM_001282927

Insert Size: 1206 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001282927.1](#)

RefSeq Size: 6122 bp

RefSeq ORF: 1206 bp

Locus ID: 10152

UniProt ID: [Q9NYB9](#)

Cytogenetics: 2q33.2

Protein Pathways: Regulation of actin cytoskeleton

MW: 44.4 kDa

Gene Summary:

Regulator of actin cytoskeleton dynamics underlying cell motility and adhesion. Functions as a component of the WAVE complex, which activates actin nucleating machinery Arp2/3 to drive lamellipodia formation (PubMed:21107423). Acts as regulator and substrate of nonreceptor tyrosine kinases ABL1 and ABL2 involved in processes linked to cell growth and differentiation. Positively regulates ABL1-mediated phosphorylation of ENAH, which is required for proper polymerization of nucleated actin filaments at the leading edge (PubMed:7590236, PubMed:8649853, PubMed:10498863). Contributes to the regulation of actin assembly at the tips of neuron projections. In particular, controls dendritic spine morphogenesis and may promote dendritic spine specification toward large mushroom-type spines known as repositories of memory in the brain (By similarity). In hippocampal neurons, may mediate actin-dependent BDNF-NTRK2 early endocytic trafficking that triggers dendrite outgrowth (By similarity). Participates in ocular lens morphogenesis, likely by regulating lamellipodia-driven adherens junction formation at the epithelial cell-secondary lens fiber interface (By similarity). Also required for nascent adherens junction assembly in epithelial cells (PubMed:15572692).[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (4) differs in its 5' UTR, differs in the presence and absence of exons in the 5' coding region, initiates translation at an alternate start codon, and lacks an in-frame exon in the central coding region, compared to variant 1. The encoded isoform (d) has a distinct N-terminus and is shorter than isoform a. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.