

Product datasheet for SC335982

Isocitrate dehydrogenase (IDH1) (NM_001282386) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Isocitrate dehydrogenase (IDH1) (NM_001282386) Human Untagged Clone
Tag:	Tag Free
Symbol:	Isocitrate dehydrogenase
Synonyms:	HEL-216; HEL-S-26; IDCD; IDH; IDP; IDPC; PICD
Vector:	pCMV6-Entry (PS100001)
Fully Sequenced ORF:	>SC335982 representing NM_001282386. Blue=Insert sequence Red=Cloning site Green=Tag(s)

```

ATGTCCAAAAAATCAGTGGCGTTCTGTGGTAGAGATGCAAGGAGATGAAATGACACGAATCATTTGG
GAATTGATTAAAGAGAACTCATTTTCCCTACGTGGAATTGGATCTACATAGCTATGATTTAGGCATA
GAGAATCGTGATGCCACCAACGACCAAGTCACCAAGGATGCTGCAGAAGCTATAAGAAGCATAATGTT
GGCGTCAAATGTCCACTATCACTCCTGATGAGAAGAGGGTTGAGGAGTTCAAGTTGAAACAAATGTGG
AAATCACCAAATGGCACCATACGAAATATTCTGGGTGGCAGGTCTTCAGAGAAGCCATTATCTGCAAA
AATATCCCCCGGCTTGTGAGTGGATGGGTAACCTATCATCATAGGTCGTCATGCTTATGGGGATCAA
TACAGAGCAACTGATTTTGTGTTCTGGGCCTGGAAAAGTAGAGATAACCTACACACCAAGTGACGGA
ACCCAAAAGGTGACATACCTGGTACATAACTTTGAAGAAGGTGGTGGTGTGCCATGGGGATGTATAAT
CAAGATAAGTCAATTGAAGATTTTGACACAGTTCCTTCCAAATGGCTCTGTCTAAGGGTTGGCCTTTG
TATCTGAGCACCAAAAACACTATTCTGAAGAAATATGATGGGCGTTTTAAAGACATCTTTCAGGAGATA
TATGACAAGCAGTACAAGTCCCAGTTTGAAGCTCAAAGATCTGGTATGAGCATAGGCTCATCGACGAC
ATGGTGGCCCAAGCTATGAAATCAGAGGGAGGCTTCATCTGGGCTGTAAAACTATGATGGTGACGTG
CAGTCGGACTCTGTGGCCCAAGGGTATGGCTCTCTCGGCATGATGACCAGCGTGCTGTTTGTCCAGAT
GGCAAGACAGTAGAAGCAGAGGCTGCCACGGGACTGTAACCCGTCACCTACCGCATGTACCAGAAAGGA
CAGGAGACGTCCACCAATCCCATTGCTTCCATTTTGCCTGGACCAGAGGGTTAGCCACAGAGCAAAG
CTTGATAACAATAAAGAGCTTGCCTTCTTTCGAAATGCTTTGGAAGAAGTCTCTATTGAGACAATTGAG
GCTGGCTTCATGACCAAGGACTTGGCTGCTTGCAATTAAGGTTTACCCAATGTGCAACGTTCTGACTAC
TTGAATACATTTGAGTTCATGGATAAACTTGGAGAAAACCTGAAGATCAAAGTCTAGCTCAGGCCAACTT
TAA
  
```

Restriction Sites:	Sgfl-MluI
ACCN:	NM_001282386
Insert Size:	1245 bp


[View online »](#)

OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001282386.1
RefSeq Size:	2313 bp
RefSeq ORF:	1245 bp
Locus ID:	3417
UniProt ID:	O75874
Cytogenetics:	2q34
Protein Pathways:	Citrate cycle (TCA cycle), Glutathione metabolism, Metabolic pathways
MW:	46.7 kDa
Gene Summary:	<p>Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydrogenase found in the cytoplasm and peroxisomes. It contains the PTS-1 peroxisomal targeting signal sequence. The presence of this enzyme in peroxisomes suggests roles in the regeneration of NADPH for intraperoxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to 3-enoyl-CoAs, as well as in peroxisomal reactions that consume 2-oxoglutarate, namely the alpha-hydroxylation of phytanic acid. The cytoplasmic enzyme serves a significant role in cytoplasmic NADPH production. Alternatively spliced transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Sep 2013]</p> <p>Transcript Variant: This variant (2) has an alternate 5' UTR exon, compared to variant 1. Variants 1, 2 and 3 encode the same protein.</p>