

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for SC336086

TADA3L (TADA3) (NM_001278270) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	TADA3L (TADA3) (NM_001278270) Human Untagged Clone
Tag:	Tag Free
Symbol:	TADA3
Synonyms:	ADA3; hADA3; NGG1; STAF54; TADA3L
Vector:	pCMV6-Entry (PS100001)
Fully Sequenced ORF:	>SC336086 representing NM_001278270. Blue=Insert sequence <mark>Red=</mark> Cloning site Green=Tag(s)
	ATGAGTGAGTTGAAAGACTGCCCCTTGCAGTTCCACGACTTCAAGTCTGTGGATCACCTGAAGGTCTGT CCCCGCTACACGGCAGTGCTGGCACGCTCTGAGGATGATGGCATCGGCATCGAGGAGCTGGACACCCTG CAGCTGGAGCTGGAGACCCTGCTGTCTTCTGCCAGCCGGCGCCTGCGTGTGCTTGAGGCCGAAACCCAG ATCCTCACCGACTGGCAGGATAAGAAAGGTGACAGACGATTCCTGAAGCTGGGTCGAGACCATGAACTT GGAGCTCCCCCCAAACATGGGAAGCCCAAGAAGCAGAAACTGGAAGGGAAGGCAGGACATGGGCCGGGC CCTGGCCCAGGACGGCCCAAATCCAAAAACCTTCAGCCCAAGAAGCAGGAAGGCAGGACATGGGCCGGGC CCTGGCCCAGGACGGCCCAAATCCAAAAACCTTCAGCCCAAGATCCAGGAATATGAATTCACTGATGAC CCTATCGACGTGCCACGGATCCCCAAAAATGATGCCCCCAACAGGTTCTGGGGCTTCAGTGGAGGCCCTAC TGTGCTGACATCACCAGCGAGGAGGTCCGCACACTTGAGGAGTTACTGAAGCCCCCAGAAGATGAGGCT GAGCATTACAAGATCCCACCCTGGGGAAGCACTACTCCCAGCGCTGGGCCCAGGAGGACCTGCTGGAG GAGCAGAAGGATGGGGCCCGGGCAGCGGCTGTGGCTGACAAGAAGAAAGGCCTCATGGGGCCCACTGACC GAACTGGACACTAAAGATGTGGATGCCCTGCTGAAGAAGTCTGAGGCCCAGCATGAACAGCCGGAAGAT GGATGCCCCTTTGGTGCCTGACGAGGCCTCCTGCAGAGGACCTGCTGGAGGAAAATATTATTTCCCCT ATGGAGGATTCTCCTATTCCTGACGAGGCCCTCCTGCAGGGCCCAGGAGGACACGGGCCAAGCACCTCCCT CGCAATCAGAACAAGCCCTTCAGTGTGCGCGCATACTAAGGAGGCCCAAGAAGAACAAGCCGGAAGAAT ATTGCCCAGGGCCTTTGGAGTCTGAGGACCGCCCCGCC
Pestriction Sites:	Sati-Miul


Restriction Sites:

Sgfl-Mlul



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Plasmid Map:



ACCN:	NM_001278270
Insert Size:	1299 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 001278270.1</u>
RefSeq Size:	1924 bp
RefSeq ORF:	1299 bp
Locus ID:	10474

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

UniProt ID:	<u>075528</u>
Cytogenetics:	3p25.3
Protein Families:	Transcription Factors
MW:	48.9 kDa
Gene Summary:	DNA-binding transcriptional activator proteins increase the rate of transcription by interacting with the transcriptional machinery bound to the basal promoter in conjunction with adaptor proteins, possibly by acetylation and destabilization of nucleosomes. The protein encoded by this gene is a transcriptional activator adaptor and a component of the histone acetyl transferase (HAT) coactivator complex which plays a crucial role in chromatin modulation and cell cycle progression. Along with the other components of the complex, this protein links transcriptional activators bound to specific promoters, to histone acetylation and the transcriptional machinery. The protein is also involved in the stabilization and activation of the p53 tumor suppressor protein that plays a role in the cellular response to DNA damage. Alternate splicing results in multiple transcript variants of this gene. [provided by RefSeq, May 2013] Transcript Variant: This variant (3) uses an alternate splice site in the 5' UTR compared to variant 1. Variants 1 and 3 encode the same isoform (a).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US