

## Product datasheet for **SC336141**

### IL31RA (NM\_001297572) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	IL31RA (NM_001297572) Human Untagged Clone
Tag:	Tag Free
Symbol:	IL31RA
Synonyms:	CRL; CRL3; GLM-R; GLMR; GPL; hGLM-R; IL-31RA; PLCA2; PRO21384
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC336141 representing NM_001297572. Blue=Insert sequence Red=Cloning site Green=Tag(s)


```
GCTCGTTTGTGAAACCGTCAGAATTTTGTAAACGACTACTATAGGGCGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGACATACTGGAGATTAGAGAATAGCGAAAACCTGAACCACCTAAGATTTTCCGTGTGAAACAGTT
TTGGGCATCAAACGAATGATTCAAATGAATGGATAAAGCCTGAGTTGGCGCCTGTTTCATCTGATTTA
AAATACACACTTCGATTCAGGACAGTCAACAGTACCAGCTGGATGGAAGTCAACTTCGCTAAGAACCGT
AAGGATAAAAACCAAACGTACAACCTCACGGGGCTGCAGCCTTTACAGAATATGTCATAGCTCTGCGA
TGTGCGGTCAAGGAGTCAAAGTTCTGGAGTGACTGGAGCCAAGAAAAATGGGAATGACTGAGGAAGAA
GCTCCATGTGGCCTGGAAGTGTGGAGAGTCTGAAACCAGCTGAGCGGATGGAAGAAGGCCAGTGCGG
TTGTTATGGAAGAAGGCAAGAGGAGCCAGCTCTAGAGAAAACACTTGGCTACAACATATGGTACTAT
CCAGAAAGCAACACTAACCTCACAGAAACAATGAACACTACTAACCAGCAGCTTGAAGTGCATCTGGGA
GGCGAGAGCTTTTGGGTGTCTATGATTTCTTATAATTCTCTTGGGAAGTCTCCAGTGGCCACCCTGAGG
ATTCCAGTATTCAAGAAAAATCATTTCAGTGCATTGAGGTCAATGCAGGCCTGCGTTGCTGAGGACCAG
CTAGTGGTGAAGTGGCAAAGCTCTGCTCTAGACGTGAACACTTGGATGATTGAATGGTTTCCGGATGTG
GACTCAGAGCCCACCACCTTCTGGGAATCTGTGTCTCAGGCCACGAAGTGGACGATCCAGCAAGAT
AAATTAACCTTTCTGGTGTATAACATCTCTGTGTATCCAATGTTGCATGACAAAGTTGGCGAGCCA
TATCCATCCAGGCTTATGCCAAAGAAGCGTTCCATCAGAAGGCTCTGAGACCAAGTGGAGAACATT
GGCGTGAAGACGGTCACGATCACATGGAAGAGATTCCCAAGAGTGAGAGAAAGGGTATCATCTGCAAC
TACACCATCTTTTACCAAGCTGAAGGTGAAAAAGGATTCTGTAAGCACGCCCATAGCGAAGTGGAAAAA
AACCCCAAGCCCCAGATAGATGCTATGGATAGACCTGTTGTAGGCATGGCTCCCCATCTCATTGTGAC
TTGCAACCTGGCATGAATCACTTAGCTTCTTAAATCTCTGAAAAATGGGGCCAAGAGCACCCACCTT
TTGGGGTTTTGGGGTTAAATGAGAGTGAAGTGACAGTACCTGAGAGGAGAGTCTGAGGAATGGAAG
GAGTTGTTATAA
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGCGC
```



[View online »](#)

Restriction Sites: SgfI-MluI

Plasmid Map:



ACCN: NM\_001297572

Insert Size: 1323 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM\\_001297572.1](#)

RefSeq Size: 2551 bp

RefSeq ORF: 1323 bp

Locus ID: 133396

UniProt ID: [Q8NI17](#)

Cytogenetics: 5q11.2

Protein Families: Druggable Genome, Transmembrane

**MW:** 50.5 kDa

**Gene Summary:** The protein encoded by this gene belongs to the type I cytokine receptor family. This receptor, with homology to gp130, is expressed on monocytes, and is involved in IL-31 signaling via activation of STAT-3 and STAT-5. It functions either as a monomer, or as part of a receptor complex with oncostatin M receptor (OSMR). Several alternatively spliced transcript variants encoding different isoforms have been noted for this gene.[provided by RefSeq, Jun 2011]

Transcript Variant: This variant (7) has multiple differences, compared to variant 1. These differences result in distinct 5' and 3' UTRs and cause translation initiation at a downstream start codon and translation termination at an alternate stop codon. The resulting isoform (7) has a shorter N-terminus and a shorter and distinct C-terminus, compared to isoform 1.