

## Product datasheet for **SC336254**

### Lipoamide Dehydrogenase (DLD) (NM\_001289752) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Lipoamide Dehydrogenase (DLD) (NM_001289752) Human Untagged Clone
Tag:	Tag Free
Symbol:	DLD
Synonyms:	DLDD; DLDH; E3; GCSL; LAD; OGDC-E3; PHE3
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

**Fully Sequenced ORF:** >SC336254 representing NM\_001289752.  
 Blue=Insert sequence Red=Cloning site Green=Tag(s)


```

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAATACGACTCACTATAGGGCGGCCGGGAATTCGTGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGCAGAGCTGGAGTCGTGTACTGCTCCTTGCCCAAGAGAGGCCATTTCAATCGAATATCTCATGGC
CTACAGGGACTTTCTGCAGTGCCTCTGAGAACTACGCAGATCAGCCGATTGATGCTGATGTAACAGTT
ATAGTTCTGGTCTGGAGGATATGTTGCTGCTATTAAAGCTGCCAGTTAGGCTTCAAGACAGTCTGC
ATTGAGAAAAATGAAACACTTGGTGGAACATGCTTGAATGTTGGTTGATTTCCTTCTAAGCCTTTATTG
AACAACTCTATTATTACCATATGGCCATGGAAAAGATTTGCATCTAGAGGAATTGAAATGTCCGAA
GTTTCGTTGAATTTAGACAAGATGATGGAGCAGAAGAGTACTGCAGTAAAAGCTTTAACAGGTGGAATT
GCCCACTATTCAAACAGAATAAGATAGATGAAGATAAATAGTGTATCTACAGGTGCTTTATCTTTA
AAAAAGTTCCAGAAAAGATGGTTGTTATTGGTGCAGGAGTAATAGGTGTAGAATTGGGTTCAAGTTG
CAAAGACTTGGTGCAGATGTGACAGCAGTTGAATTTTAGGTCATGTAGGTGGAGTTGGAATTGATATG
GAGATATCTAAAACTTTCAACGCATCCTTCAAAAACAGGGGTTAAATTTAAATTTGAATACAAAGTT
ACTGGTGTACCAAGAAGTCAGATGGAAAAATTGATGTTTCTATTGAAGCTGCTTCTGGTGGTAAAGCT
GAAGTTATCACTTGTGATGTAATCTTGGTTTGCATTGGCCGACGACCCTTACTAAGAATTTGGGACTA
GAAGAGCTGGGAATTGAACTAGATCCCAGAGGTAGAATCCAGTCAATACCAGATTTCAAATAAAATT
CCAAATATCTATGCCATTGGTGTAGTGTGCTGGTCCAATGCTGGCTCACAAGCAGAGGATGAAGGC
ATTATCTGTGTTGAAGGAATGGCTGGTGGTGTGTCACATTGACTACAATTGTGTGCCATCAGTGATT
TACACACACCTGAAGTTGCTTGGGTTGGCAAATCAGAAGAGCAGTTGAAAGAAGAGGGTATTGAGTAC
AAAGTTGGGAAATCCCATTGCTGCTAACAGCAGAGCTAAGACAAATGCTGACACAGATGGCATTGGTG
AAGATCCTTGGGCAGAAATCGACAGCAGAGACTGGGAGCACATATTCTTGGACCAGGTGCTGGAGAA
ATGGTAAATGAAGCTGCTTGGCTTGGAAATATGGAGCATCCTGTGAAGATATAGCTAGAGTCTGTGAT
GCACATCCGACCTTATCAGAAGCTTTTAGAGAAGCAAATCTTGTGCGTCATTTGGCAAATCAATCAAC
TTTTGA
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
  
```

**Restriction Sites:**

SgfI-MluI

**Plasmid Map:**



**ACCN:** NM\_001289752

**Insert Size:** 1386 bp

<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_001289752.1</a>
<b>RefSeq Size:</b>	3469 bp
<b>RefSeq ORF:</b>	1386 bp
<b>Locus ID:</b>	1738
<b>UniProt ID:</b>	<a href="#">P09622</a>
<b>Cytogenetics:</b>	7q31.1
<b>Protein Families:</b>	Druggable Genome
<b>Protein Pathways:</b>	Citrate cycle (TCA cycle), Glycine, serine and threonine metabolism, Glycolysis / Gluconeogenesis, Metabolic pathways, Pyruvate metabolism, Valine, leucine and isoleucine degradation
<b>MW:</b>	49.3 kDa
<b>Gene Summary:</b>	<p>This gene encodes a member of the class-I pyridine nucleotide-disulfide oxidoreductase family. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. In homodimeric form, the encoded protein functions as a dehydrogenase and is found in several multi-enzyme complexes that regulate energy metabolism. However, as a monomer, this protein can function as a protease. Mutations in this gene have been identified in patients with E3-deficient maple syrup urine disease and lipoamide dehydrogenase deficiency. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2014]</p> <p>Transcript Variant: This variant (4) lacks an in-frame exon in the central coding region, compared to variant 1. The encoded isoform (4) is shorter, compared to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p>