

## Product datasheet for **SC336317**

### NFAT2 (NFATC1) (NM\_001278673) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	NFAT2 (NFATC1) (NM_001278673) Human Untagged Clone
Tag:	Tag Free
Symbol:	NFATC1
Synonyms:	NF-ATC; NF-ATc1.2; NFAT2; NFATc
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

**Fully Sequenced ORF:** >SC336317 representing NM\_001278673.  
 Blue=Insert sequence Red=Cloning site Green=Tag(s)


```

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTGCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGCTGCAGCTTTTCATTGGGACGGCGGACGACCGCTGCTGCGCCCGCACGCCTTCTACCAGTGCAC
CGCATCACAGGGAAGACCGTGTCCACCACCAGCCACGAGGCCATCCTCTCCAACACCAAAGTCTGGAG
ATCCCACTCCTGCCGGAGAACAGCATGCCGAGCCGTCATTGACTGTGCCGGAATCCTGAAACTCAGAAAC
TCCGACATTGAACTTCGAAAGGAGAGACGACATCGGGAGGAAGAACACACGGGTACGGCTGGTGTTC
CGCGTTACGTCCCGCAACCCAGCGGCCGACGCTGTCCCTGCAGGTGGCCTCCAACCCCATCGAATGC
TCCCAGCGCTCAGCTCAGGAGCTGCCTCTGGTGGAGAAGCAGAGCACGGACAGCTATCCGGTCGTGGGC
GGGAAGAAGATGGTCTGTCTGGCCACAACCTCCTGCAGGACTCCAAGTCAATTTCTGGGAGAAAGCC
CCAGATGGCCACCATGTCTGGGAGATGGAAGCGAAAACGACCGGGACCTGTGCAAGCCGAATTCTCTG
GTGTTGAGATCCCGCCATTTCCGAATCAGAGGATAACCAGCCCCGTTACGTCACTTTCTACGTCTGC
AACGGGAAGAGAAGCGAAGCCAGTACCAGCGTTTCACCTACCTCCCGCCAACGTTCCAATTATAAAA
ACAGAACCCTACTGATGATTATGAGCCTGCTCCAACCTGTGGACCGGTGAGCCAGGGGTTAAGTCTCTC
CCAAGACCATACTACAGCCAGCAGCTCGCGATGCCACCCGACCCAGCTCCTGCCTCGTGCCCGGCTTC
CCGCCCTGTCCGAGAGAAGCACCTGATGCCAGCGGCCCTGGCGTGAGCCCCAAGCTCCACGACCTT
TCTCCCGTGCCTACACCAAGGGCGTTGCCAGCCCGGGCCACTGTACCTCGGACTCCCGCAGCCGGCC
GGAGAGGCCCCCGCGTCCAGGACGTGCCAGGCCAGTGGCCACGCACCCCGGCTCGCCCGGGCAGCCA
CCCCGGCCCTGTGCCACAGCAGGTGAGTGGCCCTCAAGCAGTAGTGCCCCCTGGTCTCGAACAC
TCGCTGTCCCCAGCAGCCCTCTCTCCACTCCCGCTGCCACCAAGAGCCGACCTCGCTGCAGCCC
TGCAGCCACGCGTGCCCGCCCGCCAGGGCCCGCCAGCACCTGCCGTCCACGCTCCGAGGACGAG
TCTCCGACTGCCGGGCCACGGCTGCTGCCAGAGGTGCATGAGGACGGTAGTCTAATTTGGCCCTATT
CCTGTAACGCTAAGCGAGGCCTGAAGAGTTGGACCAGTTGTACCTGGATGACGTAATGAAATAATA
CGAAATGACCTCTCCAGCACGAGCACCCACTCTAG
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
  
```

**Restriction Sites:**

SgfI-MluI

**Plasmid Map:**



**ACCN:** NM\_001278673

**Insert Size:** 1416 bp

<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_001278673.1</a>
<b>RefSeq Size:</b>	3558 bp
<b>RefSeq ORF:</b>	1416 bp
<b>Locus ID:</b>	4772
<b>UniProt ID:</b>	<a href="#">O95644</a>
<b>Cytogenetics:</b>	18q23
<b>Protein Families:</b>	Druggable Genome, Transcription Factors
<b>Protein Pathways:</b>	Axon guidance, B cell receptor signaling pathway, Natural killer cell mediated cytotoxicity, T cell receptor signaling pathway, VEGF signaling pathway, Wnt signaling pathway
<b>MW:</b>	51.1 kDa
<b>Gene Summary:</b>	<p>The product of this gene is a component of the nuclear factor of activated T cells DNA-binding transcription complex. This complex consists of at least two components: a preexisting cytosolic component that translocates to the nucleus upon T cell receptor (TCR) stimulation, and an inducible nuclear component. Proteins belonging to this family of transcription factors play a central role in inducible gene transcription during immune response. The product of this gene is an inducible nuclear component. It functions as a major molecular target for the immunosuppressive drugs such as cyclosporin A. Multiple alternatively spliced transcript variants encoding distinct isoforms have been identified for this gene. Different isoforms of this protein may regulate inducible expression of different cytokine genes. [provided by RefSeq, Jul 2013]</p> <p>Transcript Variant: This variant (10) has an alternate 5' terminal exon and lacks an internal exon in the 5' region, which results in a downstream translation start codon, compared to variant 6. The encoded isoform (J) is shorter at the N-terminus, compared to isoform F.</p>