

Product datasheet for **SC336389**

CD30 (TNFRSF8) (NM_001281430) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	CD30 (TNFRSF8) (NM_001281430) Human Untagged Clone
Tag:	Tag Free
Symbol:	CD30
Synonyms:	CD30; D1S166E; Ki-1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >SC336389 representing NM_001281430.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)

```

GCTCGTTTAGTGAACCGTCAGAATTTTGAATACGACTCACTATAGGGCGGCCGGGAATTCGTGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGTTCTGTTCACGTCCTGCCGTCCAACTCCTGTGCCCGCTGCTTCTTCCATTCTGTCTGTCCGGCAGGG
ATGATTGTCAAGTTCCCAGGCACGGCGCAGAAGAACCGTCTGTGAGCCGGCTTCCCAGGGGTACGC
CCTGCCTGTGCCAGCCCAGAGAACTGCAAGGAACCCTCCAGTGGCACCATCCCCAGGCCAAGCCCACC
CCGGTGTCCCAGCAACCTCCAGTGCCAGCACCATGCCTGTAAGAGGGGGCACCCGCCTCGCCCAGGAA
GCTGTTCTAAACTGACGAGGGCTCCCGACTCTCCCTCCTCTGTGGGAAGGCCTAGTTCCAGATCCAGGT
CTGTCCCCAACACAGCCATGCCAGAGGGGTCTGGTATTGCAGAAAGCAGTGTGAGCCCGACTACTAC
CTGGACGAGGCCCGCTGCACGGCTCGTGAGCTGTTCTCGAGATGACCTTGTGGAGAAGACGCCA
TGTGCATGGAACTCCTCCGCACCTGCGAATGTCGACCTGGCATGATCTGTGCCACATCAGCCACCAAC
TCCTGTGCCCGCTGTGTCCCTACCCAATCTGTGCAGCAGAGACGGTCACCAAGCCCCAGGATATGGCT
GAGAAGGACACCACCTTTGAGGGCCACCCTGGGGACCCAGCCGACTGCAACCCACCCAGAGAAT
GGCGAGGGCGCTGCCAGCACAGCCCACTCAGAGCTTGTGGTGGACTCCCAGGCCAGTAAAGACGCTG
CCCATCCCAACCAGCGCTCCCGTCGCTCTCCTCCACGGGGAAGCCCGTTCTGGATGCAGGGCCAGTG
CTTTCTGGGTGATCCTGGTGTGGTGTGGTGGTCCAGCGCCTTCTCCTGTGCCACCGGAGG
GCCTGCAGGAAGCGAATTCGGCAGAAGCTCCACCTGTGCTACCCGGTCCAGACCTCCCAGCCAAAGCTA
GAGCTTGTGGATTCCAGACCCAGGAGGAGCTCAACGCTGAGGAGTGGTGCCTCGGTGACAGAACCCGTC
GCGGAAGAGCGAGGGTTAATGAGCCAGCCACTGATGGAGACCTGCCACAGCGTGGGGGCAGCCTACCTG
GAGAGCCTGCCGCTGCAGGATGCCAGCCCGCCGGGGCCCTCGTCCCCAGGGACCTTCTGAGCCC
CGGGTGTCCACGGAGCACACCAATAACAAGATTGAGAAAATCTACATCATGAAGGCTGACACCGTGATC
GTGGGGACCGTGAAGGCTGAGCTGCCGGAGGGCCGGGGCCTGGCGGGGCCAGCAGAGCCCGAGTTGGAG
GAGGAGCTGGAGGCGGACCATACCCCCACTACCCCGAGCAGGAGACAGAACCGCCTCTGGGCAGCTGC
AGCGATGTATGCTCTCAGTGGAAAGAGGAAAGAACCCCTTGCCCACAGCTGCCTCTGAAAG
TGA
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
  
```

- Restriction Sites:** SgfI-MluI
- ACCN:** NM_001281430
- Insert Size:** 1452 bp
- OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
- Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
- Reconstitution Method:**
1. Centrifuge at 5,000xg for 5min.
 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
 3. Close the tube and incubate for 10 minutes at room temperature.
 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq:	NM_001281430.2
RefSeq Size:	3616 bp
RefSeq ORF:	1452 bp
Locus ID:	943
UniProt ID:	P28908
Cytogenetics:	1p36.22
Protein Families:	Druggable Genome, ES Cell Differentiation/IPS, Stem cell - Pluripotency, Transmembrane
Protein Pathways:	Cytokine-cytokine receptor interaction
MW:	50.9 kDa
Gene Summary:	<p>The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is expressed by activated, but not by resting, T and B cells. TRAF2 and TRAF5 can interact with this receptor, and mediate the signal transduction that leads to the activation of NF-kappaB. This receptor is a positive regulator of apoptosis, and also has been shown to limit the proliferative potential of autoreactive CD8 effector T cells and protect the body against autoimmunity. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]</p> <p>Transcript Variant: This variant (3) has multiple differences, compared to variant 1. These differences result in a distinct 5' UTR and cause translation initiation at a downstream start codon, compared to variant 1. The encoded protein (isoform 3) is shorter than isoform 1.</p>