

Product datasheet for **SC336432**

COP1 (RFWD2) (NM_001286644) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	COP1 (RFWD2) (NM_001286644) Human Untagged Clone
Tag:	Tag Free
Symbol:	COP1
Synonyms:	CFAP78; FAP78; RFWD2; RNF200
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >SC336432 representing NM_001286644.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)


```

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTGACTG
GATCCGGTACCGAGGAGATCTGCCGCCCGGATCGCC
ATGTTGTGGACAATATTGACCATCTGTATCCTAATTTCTTGGGAATCACATGCAGCCCAACTACAGATT
CTTATGGAAATTCCTCAAGTTGCAAGAAGAAATAGAGAGAGCAACTGGAACAGATCCAGAAGGAGCTA
AGTGTTTTGGAAGAGGATATTAAGAGAGTGGAAGAAATGAGTGGCTTATACTCTCCTGTGAGGAT
AGCACAGTGCCTCAATTTGAAGCTCCTTCTCCATCACACAGTAGTATTATTGATTCCACAGAATACAGC
CAACCTCCAGTTTCAGTGGCAGTTCTCAGACAAAGAACAGCCTTGGTATAATAGCACGTTAGCATCA
AGACGAAAACGACTTACTGCTCATTTTGAAGACTTGGAGCAGTGTACTTTTCTACAAGGATGTCTCGT
ATCTCAGATGACAGTCAACTGCAAGCCAGTTGGATGAATTTAGGAATGCTTGTCCAAGTTTACTCGA
TATAATTCAGTACGACCTTAGCCACATTGTCATATGCTAGTGTCTCTATAATGGTTCAGTATAGTC
TCTAGTATTGAATTTGACCGGGATTGTGACTATTTTGGCATTGCTGGAGTTACAAAGAAGATTAAGTC
TATGAATATGACTGTATCCAGGATGCAGTGGATATTCATTACCTGAGAATGAAATGACCTGCAAT
TCGAAAATCAGCTGTATCAGTTGGAGTAGTACCATAAGAACCTGTTAGCTAGCAGTGATTATGAAGGC
ACTGTTATTTTATGGGATGGATTACAGGACAGAGGTCAAAGGTCTATCAGGAGCATGAGAAGAGGTGT
TGGAGTGTGACTTTAATTTGATGGATCCTAAACTCTTGGCTTCAGGTTCTGATGATGCAAAAGTGAAG
CTGTGGTCTACCAATCTAGACAACCTCAGTGGCAAGCATTGAGGCAAAAGGCTAATGTGTGCTGTGTTAAA
TTCAGCCCCTCTCCAGATACCATTTGGCTTTCGGCTGTGCAGATCACTGTGCCACTACTATGATCTT
CGTAACACTAAACAGCCAATCATGGTATTCAAAGGACACCGTAAAGCAGTCTCTTATGCAAAGTTTGTG
AGTGGTGAGGAAATTGTCTCTGCCTCAACAGACAGTCACTAAAACCTGTGGAATGTAGGGAACCATAC
TGCCACGTTCCCTCAAGGGTCATATCAATGAAAAAACTTTGTAGCCCTGGCTTCCAATGGAGATTAT
ATAGCTTGTGGAAGTAAAATAACTCTCTACTGTACTATAAAGGACTTTCTAAGACTTTGCTAACT
TTTAAGTTTGATACAGTCAAAGTGTCTCGACAAAGACCGAAAAGAAGATGATACAAATGAATTTGTT
AGTGTGTGTGCTGGAGGGCACTACCAGATGGGGAGTCCAATGTGCTGATTGCTGCTAACAGTCAGGGT
ACAATTAAGGTGCTAGAATTGGTATGA
ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
  
```

Restriction Sites:

SgfI-MluI

Plasmid Map:



ACCN:

NM_001286644

Insert Size:	1476 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001286644.1
RefSeq Size:	2609 bp
RefSeq ORF:	1476 bp
Locus ID:	64326
UniProt ID:	Q8NHY2
Cytogenetics:	1q25.1-q25.2
Protein Pathways:	p53 signaling pathway, Ubiquitin mediated proteolysis
MW:	55.7 kDa
Gene Summary:	<p>E3 ubiquitin-protein ligase that mediates ubiquitination and subsequent proteasomal degradation of target proteins. E3 ubiquitin ligases accept ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates. Involved in JUN ubiquitination and degradation. Directly involved in p53 (TP53) ubiquitination and degradation, thereby abolishing p53-dependent transcription and apoptosis. Ubiquitinates p53 independently of MDM2 or RCHY1. Probably mediates E3 ubiquitin ligase activity by functioning as the essential RING domain subunit of larger E3 complexes. In contrast, it does not constitute the catalytic RING subunit in the DCX DET1-COP1 complex that negatively regulates JUN, the ubiquitin ligase activity being mediated by RBX1. Involved in 14-3-3 protein sigma/SFN ubiquitination and proteasomal degradation, leading to AKT activation and promotion of cell survival. Ubiquitinates MTA1 leading to its proteasomal degradation. Upon binding to TRIB1, ubiquitinates CEBPA, which lacks a canonical COP1-binding motif (Probable).[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (3) differs in its 5' UTR and initiates translation at an alternate start codon, compared to variant 1. The encoded isoform (c) has a shorter and distinct N-terminus, compared to isoform a.</p>