

Product datasheet for **SC336492**

NADPH oxidase 4 (NOX4) (NM_001291926) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	NADPH oxidase 4 (NOX4) (NM_001291926) Human Untagged Clone
Tag:	Tag Free
Symbol:	NOX4
Synonyms:	KOX; KOX-1; RENOX
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >SC336492 representing NM_001291926.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)


```

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTGCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCCGGATCGCC
ATGTGCCGAACACTCTTGGCTTACCTCCGAGGATCACAGAAGTTCCAAGCAGGAGAACCAGGAGATTG
TTGGATAAAAGCAGAACATTCCATATTACCTGTGGTGTACTATCTGTATTTTCTCAGCGTGCATGTG
GCTGCCCATCTGGTGAATGCCCTCAACTTCTCAGTGAATTACAGTGAAGACTTTGTTGAAGTGAATGCA
GCAAGATACCGAGATGAGGATCCTAGAAAACTTCTCTTCACAACTGTTCCCTGGCCTGACAGGGGTCTGC
ATGGTGGTGGTGTATTCTCATGATCACAGCCTCTACATATGCAATAAGAGTTTCTAACTATGATATC
TTCTGGTATACTATAACCTCTTCTTTGTCTTCTACATGCTGCTGACGTTGCATGTTTCAGGAGGGCTG
CTGAAGTATAAACTAATTTAGATACCCACCCTCCCGGCTGCATCAGTCTTAACCGAACAGCTCTCAG
AATATTTCTTACCAGAGATTTTCTCAGAACATTTTCATGAACCTTTCCCTGAAGGATTTTCAAACCG
GCAGAGTTTACCAGCACAAATTTGTGAAGATTTGTATGGAAGAGCCAGATTCCAAGCTAATTTCCA
CAGACTTGGCTTTGGATTTCTGGACCTTTGTGCCTGTACTGTGCCGAAAGACTTTACAGGTATATCCGG
AGCAATAAGCCAGTCACCATCATTTCCGGTCATAAGTCATCCCTCAGATGTCATGGAATCCGAATGGTC
AAAGAAAATTTAAAGCAAGACCTGGTCAGTATATTACTCTACATTGTCCAGTGTATCTGCATTAGAA
AATCATCCATTTACCCTACAATGTGTCCAACCTGAAACCAAAGCAACATTTGGGGTTTCATCTTAAATA
GTAGGAGACTGGACAGAACGATTTCCGAGATTTACTACTGCCTCCATCTAGTCAAGACTCCGAAATTTCTG
CCCTTCATTCAATCTAGAAAATATCCCAAGCTGTATATTGATGGTCCTTTTGGAAGTCCATTTGAGGAA
TCACTGAACTATGAGGTCAGCCTCTGCGTGGCTGGAGGCATTGGAGTAACTCCATTTGCATCAATACTC
AACACCCTGTTGGATGACTGAAACCATAACAAGCTTAGAAGACTATACTTTATTTGGGTATGCAGAGAT
ATCCAGTCCCTCCGTTGGTTGCGAGATTTACTCTGTATGTTGCATAACAAGTTTTGGCAAGAGAACAGA
CCTGACTATGTCAACATCCAGCTGTACCTCAGTCAAACAGATGGGATACAGAAGATAATTGGAGAAAAA
TATCATGCACTGAATTAAGACTGTTTATAGGACGTCCTCGGTGGAACTTTTGTGGTGAATAGCA
AAATATAACAGAGGAAAAACAGTTGGTGTCTTCTGTTGTGGACCAATTCATATCCAAGACTCTTCAT
AAACTGAGTAACCGAACAACCTCATATGGGACAAGATTTGAATACAATAAAGAGTCTTTCCAGTGA
ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
  
```

Restriction Sites:

SgfI-MluI

Plasmid Map:



ACCN:

NM_001291926

Insert Size:	1515 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001291926.1
RefSeq Size:	4455 bp
RefSeq ORF:	1515 bp
Locus ID:	50507
UniProt ID:	Q9NPH5
Cytogenetics:	11q14.3
Protein Families:	Druggable Genome, Transmembrane
MW:	58.4 kDa
Gene Summary:	<p>This gene encodes a member of the NOX-family of enzymes that functions as the catalytic subunit the NADPH oxidase complex. The encoded protein is localized to non-phagocytic cells where it acts as an oxygen sensor and catalyzes the reduction of molecular oxygen to various reactive oxygen species (ROS). The ROS generated by this protein have been implicated in numerous biological functions including signal transduction, cell differentiation and tumor cell growth. A pseudogene has been identified on the other arm of chromosome 11. Alternative splicing results in multiple transcript variants.[provided by RefSeq, Jan 2009]</p> <p>Transcript Variant: This variant (4) differs in its 5' UTR, and uses a downstream start codon, compared to variant 1. The encoded isoform (d) has a shorter N-terminus, compared to isoform a. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p>