

Product datasheet for **SC336532**

DCP1A (NM_001290205) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	DCP1A (NM_001290205) Human Untagged Clone
Tag:	Tag Free
Symbol:	DCP1A
Synonyms:	HSA275986; Nbla00360; SMAD4IP1; SMIF
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)


Fully Sequenced ORF: >SC336532 representing NM_001290205.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)

```

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTGCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCACGATCGCC
ATGAACCATTTCTTCTGTATAGAATGCAAGCTGTTGAGAAAACAAGAGGCTCAGAGCTGTTTAGGAAT
CTGCTCGAGATTCGAAAGTGTGATATATAGTATCTGGTTTTATGACAAGAATGACTGTCACCGCATA
GCAAAACTCATGGCTGATGTGGTAGAAGAGGAGACACGGCGATCCCAGCAAGCTGCTCGGGACAAACAG
AGTCCCAGCCAGGCCAATGGCTGCACGACCACAGGCCCATCGACATCCTGGAGATGCTGAGCAGAGCC
AAGGATGAGTATGAGAGGAATCAGATGGGTGACTCAAATATCTCCAGCCCTGGGTACAGCCAAGCACT
CAGCTCTCCAATCTGGGAAGCACCGAGACTCTAGAAGAAATGCCCTCCGGGTACAGGATAAGTCTGCT
CCATCTGGACACAAGCATCTGACGGTAGAAGAGTTATTTGGAACCTTTTGCCAAAGGAACAACCAGCA
GTTGTGGTCTGGATTGAGAAGAAATGGAGAGTTGCCAGGAGATGCCTCCAGAAAGAGCCCAATTCA
TTCTACCATTTCCCTTTGAGCAGTTAGGAGGAGCCCTCAATCAGAAACCTGGGTGTCCTTCTGCT
GCCACCATTGAGTCCAGCTGAAATCACACCCCGGTGCTAATCACTCCAGCCTCCATCACACAGTCC
AATGAAAAGCATGCTCCAACCTACACAATCCCCTTGAGCCCTGTTCTCAGTCCCCTCTGCCAGTGAA
GCTCCTACTGCACAGGTTCCCCCAGCTTACCTCGAAACAGCACCATGATGCAGGCAAGTGAAGACCAG
CCTAGACAGAGGTCTCCACTCCTGAACCAGCCAGTCCCTGAGCTAAGCCATGCCAGTCTGATTGCCAAC
CAGAGCCCTTCAGGGCCCCATTGAACGTGACGAACACAGCTGGCACATCCCTCCAAGCGTTGATCTT
CTCCAGAAACTCAGGTTGACCCACAGCATGACCAAATACAGACACAACCACTTGGGAAAGGTGCAATG
GTAGCCAGCTTTTCTCCGGCAGCTGGTCAGCTAGCCACACCTGAGAGCTTCATAGAGCCTCCCTCTAAG
ACAGCAGCAGCAAGAGTGGCGCCTCAGCTCCCTGAGCAACATGGTGTGCTCCCTTCAGTCTATG
CAGCAGAACCAGGATCCTGAAGTATTTGTGCAGCCTAAGGTGTTATCCAGTGCCATCCCGGTTGCAGGC
GCCCACTGGTTACTGCAACGACCACTGAGTGTCTTCAGTCTGCTGGCCCAAGTGTTTTCCAGCAG
ACAGTTACAAGATCTTCGACCTTGAGAGGAAAGCCAGTCCCCTTCTCCTTAACCTATTGGAACGCCA
GAAAGTCAGAGAAAGCCTTCCATTATTCTCAGCAAGTCTCAGCTCCAGGATACATTAATACATCTAATA
AAGAATGATTCCAGTTCCTCAGTACACTTCATGAAGTCTACTTGAGGTTCTGACCAAGAACAAGAC
AACCACAACCTATGA
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTAAACGGCCGGC
  
```

Restriction Sites: Sgfl-MluI

Plasmid Map:



ACCN: NM_001290205

Insert Size:	1533 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001290205.1
RefSeq Size:	6082 bp
RefSeq ORF:	1533 bp
Locus ID:	55802
UniProt ID:	Q9NPI6
Cytogenetics:	3p21.1
Protein Families:	Transcription Factors
Protein Pathways:	RNA degradation
MW:	55.2 kDa
Gene Summary:	<p>Decapping is a key step in general and regulated mRNA decay. The protein encoded by this gene is a decapping enzyme. This protein and another decapping enzyme form a decapping complex, which interacts with the nonsense-mediated decay factor hUpf1 and may be recruited to mRNAs containing premature termination codons. This protein also participates in the TGF-beta signaling pathway. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Feb 2014]</p> <p>Transcript Variant: This variant (3) contains an additional internal exon in the 5' region, resulting in translation initiation at an alternate start codon, compared to variant 1. The encoded isoform (c) has a distinct N-terminus and is shorter than isoform a. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p>