

Product datasheet for **SC336560**

NISCH (NM_001276294) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	NISCH (NM_001276294) Human Untagged Clone
Tag:	Tag Free
Symbol:	NISCH
Synonyms:	hIRAS; I-1; IR1; IRAS
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >SC336560 representing NM_001276294.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)


```

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTGCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCACGATCGCC
ATGGCGACCGCGCACCTTCGGGCCAGCGGGAAGCCGAGCCGGCCAAGGAAGCGCGCGTCTGTTGGC
TCGGAGCTTGTGGACACTATACGGTTTACATCATCCAGGTCAGTGCAGCCATGAGTGGACAGTA
AAGCACCCTACAGCGACTTCCATGACCTGCATGAAAAGCTCGTTGCAGAGAGAAAGATTGATAAAAAT
CTGCTTCCGCCAAAAAGATAATTGGGAAAACTCAAGAAGCTTGGTGGAGAAGAGGGAGAAGGATCTG
GAGGTCTACCTCCAGAAGCTCCTGGCTGCCTCCCTGGCGTGACCCCGAGTACTGGCCCACTTCTTG
CATTTTCACTTCTATGAGATAAATGGCATCACCGCGCACTGGCTGAAGAGCTCTTTGAGAAAGGAGAA
CAGCTCCTGGGGCCGGCGAGGTCTTTGCCATTGGACCCCTGCAGCTGTATGCCGTACGGAGCAGCTG
CAGCAGGAAAGCCACGTGCGCCAGTGGGGATGCCAAGACCGACCTCGGGCACATCCTGGACTTACC
TGTCGCCTTAAGTACCTAAGGTTTCTGGCACAGAAGGACCTTTTGGACAGCAACATTCAGGAGCAG
CTCCTGCCGTTTCGACCTATCAATATTCAAGTCCCTGCATCAGGTGGAGATAAGTCACTGTGATGCTAAG
CACATCAGAGGGCTGGTGCATCGAAGCCACCTTAGCCACGCTGAGTGTCCGCTTCTCAGCAACCTCG
ATGAAGGAAGTCCTTGTTCCTGAAGCCTCAGAATTTGATGAGTGGGAGCCTGAAGGCACAACCTAGAA
GGCCCTGTGACTGCCGTATCCCACTTGGCAGGCATTGACCACGCTTGACCTGAGCCACAACAGCATC
TCCGAGATCGACGAGTCTGTGAACTGATCCCAAAGATTGAGTTCCTGGACCTGAGTCACAATGGATTG
CTGTTTGTGGACAATCTGCAGCACCTGTATAACCTTGTGCATCTGGACCTGTCTACAACAAGCTCTCC
TCCTTGAAGGGCTTACACCAAGCTGGGAACATCAAGACCTTAAACCTGGCAGCAACCTCCTAGAG
AGTCTGAGTGGCTGCACAAGCTTACTCACTGGTCAACCTGGATCTCCGGGACAACAGGATCGAACAG
ATGGAGGAGTCCGGAGCATAGGCAGCCTCCCGTGTCTGGAGCAGTGTCTCTGCTGAACAACCTCTG
AGCATCATCCCCGACTACCGGACCAAGGTGCTGGCTCAGTTCGGAGAGAGGGCCTCAGAGGTCTGTCTG
GATGACACAGTGACCACAGAGAAGGAGCTGGACACTGTGGAAGTCTGAAAGCAATTGAGAAAGCCAAG
GAGGTCAAGTCCAACTGAGCAACCCAGAGAAGAAGGGTGGTGAAGACTCCCGGCTCTCAGCTGCCCCC
TGCATCAGACCCAGCAGCTCCCCTCCCCTGCTGGCTCCCGCATCTGCCTCCCTGCCCCAGCCATCCTC
TCTAACCAAGGCATCCTCGGAGATGAGTGA
ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
  
```

Restriction Sites:

SgfI-MluI

Plasmid Map:



ACCN:

NM_001276294

Insert Size:	1548 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001276294.1</u>
RefSeq Size:	2706 bp
RefSeq ORF:	1548 bp
Locus ID:	11188
UniProt ID:	<u>Q9Y2I1</u>
Cytogenetics:	3p21.1
Protein Families:	Druggable Genome
MW:	56.9 kDa

Gene Summary:

This gene encodes a nonadrenergic imidazoline-1 receptor protein that localizes to the cytosol and anchors to the inner layer of the plasma membrane. The orthologous mouse protein has been shown to influence cytoskeletal organization and cell migration by binding to alpha-5-beta-1 integrin. In humans, this protein has been shown to bind to the adapter insulin receptor substrate 4 (IRS4) to mediate translocation of alpha-5 integrin from the cell membrane to endosomes. Expression of this protein was reduced in human breast cancers while its overexpression reduced tumor growth and metastasis; possibly by limiting the expression of alpha-5 integrin. In human cardiac tissue, this gene was found to affect cell growth and death while in neural tissue it affected neuronal growth and differentiation. Alternative splicing results in multiple transcript variants encoding different isoforms. Some isoforms lack the expected C-terminal domains of a functional imidazoline receptor. [provided by RefSeq, Jan 2013]

Transcript Variant: This variant (3) lacks several exons and includes an alternate 3' terminal exon, resulting in a novel 3' coding region and 3' UTR, compared to variant 1. It encodes isoform 3 (also known as IRAS-S) which is shorter and has a distinct C-terminus, compared to isoform 1. This isoform lacks multiple domains which are believed to be required by functional imidazoline receptors. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.