

Product datasheet for **SC336571**

CDC25B (NM_001287516) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	CDC25B (NM_001287516) Human Untagged Clone
Tag:	Tag Free
Symbol:	CDC25B
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)


Fully Sequenced ORF: >SC336571 representing NM_001287516.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)

```

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTGCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGCACAGCGAAACCCCAAGAGTCAGGTAGGGACCCCTGCTCTTCCGAGCCGACGCCCCCTGACGCAC
CTATCCCTGTCTCGACGGGCATCCGAATCCTCCCTGCTGCTGAATCCTCCGAATCTTCTGATGCAGGT
CTCTGCATGGATTCCCCCAGCCCTATGGACCCCCACATGGCGGAGCAGAGTTTGAACAGGCCATCCAG
GCAGCCAGCCGGATCATTGAAACGAGCAGTTTGCATCAGACGCTTCCAGTCTATGCCGGTGAGGCTG
CTGGGCCACAGCCCGTGCTTCGGAACATCACAACTCCCAGGCGCCGACGCGCCGGAGGAAGAGCGAG
GCGGGCAGTGGAGCTGCCAGCAGCTCTGGGAAGACAAGGAGAATGATGGATTTGCTTCAAGATGCCA
TGAAGCCACACATCCCAGCTCCACCCATGCTCTGCCAGAGTGGCCAGCCGACGGGAAGCCTTTGCC
CAGAGACCCAGCTCGGCCCGACCTGATGTGTCTCAGTCTGACCGAAGATGGAAGTGGAGGAGCTC
AGCCCCCTGGCCCTAGGTCGCTTCTCTGACCCCTGCAGAGGGGATACTGAGGAAGATGATGGATTT
GTGGACATCCTAGAGAGTGACTTAAAGGATGATGATGCAGTTCGCCAGGCATGGAGAGTCTCATTAGT
GCCCCACTGGTCAAGACCTTGGAAAAGGAAGAGGAAAAGGACCTCGTCATGTACAGCAAGTGCCAGCGG
CTTCCGCTCTCCGTCCATGCCCTGCAGCGTGATCCGGCCCATCCTCAAGAGGCTGGAGCGGCCCCAG
GACAGGGACACGCCCGTGCAATAAGCGGAGGCGGAGCGTGACCCCTCCTGAGGAGCAGCAGGAGGCT
GAGGAACCTAAAGCCCGCTCCTCCGCTCAAAATCACTGTGCACGATGAGATCGAGAACCTCCTGGAC
AGTGACCACCGAGAGCTGATTGGAGATTACTCTAAGGCCCTTCTCCTACAGACAGTAGACGGAAAGCAC
CAAGACCTCAAGTACATCTACCAGAAACGATGGTGGCCCTATTGACGGGCAAGTTCAGCAACATCGTG
GATAAGTTTGTGATTGTAGACTGCAGATACCCCTATGAATATGAAGGCGGGCACATCAAGACTGCGGTG
AACTTGCCCTGGAACGCGAGCCGAGAGCTTCTACTGAAGAGCCCATCGGCCCTGTAGCCTGGAC
AAGAGAGTCATCCTCATTTCCTACTGTGAATTCTCATCTGAGCGTGGGCCCGCATGTGCCGTTTATC
AGGAACGAGACCGTGCTCAACGACTACCCAGCCTCTACTACCCTGAGATGTATATCCTGAAAGGC
GGCTACAAGGAGTTCTTCCCTCAGCACCCGAACTTCTGTGAACCCAGGACTACCGGCCATGAACCAC
GAGGCCTCAAGGATGAGCTAAAGACCTTCCGCTCAAGACTCGCAGCTGGGCTGGGAGCGGAGCCGG
CGGGAGCTCTGTAGCCGGCTGCAGGACCAGTGA
ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
  
```

Restriction Sites: SgfI-MluI

Plasmid Map:



ACCN: NM_001287516

Insert Size:	1551 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001287516.1</u>
RefSeq Size:	3071 bp
RefSeq ORF:	1551 bp
Locus ID:	994
Cytogenetics:	20p13
Protein Families:	Druggable Genome, Phosphatase
Protein Pathways:	Cell cycle, MAPK signaling pathway, Progesterone-mediated oocyte maturation
MW:	58.9 kDa
Gene Summary:	<p>CDC25B is a member of the CDC25 family of phosphatases. CDC25B activates the cyclin dependent kinase CDC2 by removing two phosphate groups and it is required for entry into mitosis. CDC25B shuttles between the nucleus and the cytoplasm due to nuclear localization and nuclear export signals. The protein is nuclear in the M and G1 phases of the cell cycle and moves to the cytoplasm during S and G2. CDC25B has oncogenic properties, although its role in tumor formation has not been determined. Multiple transcript variants for this gene exist. [provided by RefSeq, Jul 2008]</p> <p>Transcript Variant: This variant (4) differs in the 5' UTR and initiates translation at an upstream start codon, compared to variant 1. The resulting protein (isoform 4) has a shorter N-terminus compared to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p>