

## Product datasheet for **SC336635**

### PRMT5 (NM\_001282954) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	PRMT5 (NM_001282954) Human Untagged Clone
Tag:	Tag Free
Symbol:	PRMT5
Synonyms:	HRMT1L5; HSL7; IBP72; JBP1; SKB1; SKB1Hs
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

**Fully Sequenced ORF:** >SC336635 representing NM\_001282954.  
 Blue=Insert sequence Red=Cloning site Green=Tag(s)


```

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTGACTG
GATCCGGTACCGAGGAGATCTGCCGCCCGGATCGCC
ATGTTACAGGAGCTGAATTTTGGTGCATATTTGGGTCTTCCAGCTTTCCTGCTGCCCTTAATCAGGAA
GATAACACCAACCTGGCCAGAGTTTGGACCAACCACACACTGCCATCACTCTCCATGTTCTGG
ATGCGGGTACCCTTGGTGGCACCAGAGGACCTGAGAGATGATATAATTGAGAATGCACCAACTACACAC
ACAGAGGAGTACAGTGGGAGGAGAAAACGTGGATGTGGTGGCACAACCTCCGGACTTGTGTGACTAT
AGTAAGAGGATTGCAGTGGCTCTTGAATTGGGGCTGACCTCCCATCTAATCATGTCATTGATCGCTGG
CTTGGGGAGCCATCAAAGCAGCCATTCTCCCACTAGCATTTCCTGACCAATAAGAAGGGATTTCCT
GTTCTTTCTAAGATGCACCAGAGGCTCATCTCCGGCTCCTCAAGTTGGAGGTGCAGTTCATCATACA
GGCACCAACCACCACTCAGAGAAGGAGTCTGCTCCTACCTCCAATACCTGGAATACTTAAGCCAGAAC
CGTCTCCACCTAATGCCTATGAACTCTTGGCAAGGGCTATGAAGACTATCTGCAGTCCCCGCTTCAG
CCACTGATGGACAATCTGGAATCTCAGACATATGAAGTGTGAAAAGGACCCCATCAAATACTCTCAG
TACCAGCAGGCCATCTATAAATGTCTGCTAGACCGAGTACCAGAAGAGGAGAAGGATACCAATGTCCAG
GTACTGATGGTGTGGGAGCAGGACGGGACCCCTGGTGAACGCTTCCCTGCGGGCAGCCAAGCAGGCC
GACCGGGGATAAAGCTGTATGCTGTGGAGAAAAACCAATGCCGTGGTACGCTAGAGAACTGGCAG
TTTGAAGAATGGGAAGCCAAGTGACCGTAGTCTCATCAGACATGAGGGAATGGTGGCTCCAGAGAAA
GCAGACATCATTGTCAGTGAGCTTCTGGGCTCATTGCTGACAATGAATTGTCGCTGAGTGCCTGGAT
GGAGCCCAGCACTTCTAAAAGATGATGGTGTGAGCATCCCCGGGAGTACACTTCTTTCTGGCTCCC
ATCTCTTCTCCAAGCTGTACAATGAGGTCCGAGCCTGTAGGGAGAAGGACCGTGACCCCTGAGGCCAG
TTTGAGATGCCTTATGTGGTACGGCTGCACAACCTCCACCAGCTCTCTGCACCCAGCCCTGTTTACC
TTCAGCCATCCCAACAGAGATCCTATGATTGACAACAACCGCTATTGCACCTTGGAAATTTCTGTGGAG
GTGAACACAGTACTACATGGCTTTGCCGCTACTTTGAGACTGTCTTTATCAGGACATCACTCTGAGT
ATCCGTCCAGAGACTCACTCTCCTGGGATGTTCTCATGGTTTCCCATCTCTTCCCTATTAAGCAGCCC
ATAACGGTACGTGAAGGCCAAACCATCTGTGTGCGTTTCTGGCGATGCAGCAATTCCAAGAAGGTGTGG
TATGAGTGGGCTGTGACAGCACCAGTCTGTTCTGCTATTCATAACCCACAGGCCGCTCATATACCATT
GGCCTTAG
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGCGC
  
```

**Restriction Sites:**

SgfI-MluI

**Plasmid Map:**



<b>ACCN:</b>	NM_001282954
<b>Insert Size:</b>	1596 bp
<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_001282954.1</a>
<b>RefSeq Size:</b>	2507 bp
<b>RefSeq ORF:</b>	1596 bp
<b>Locus ID:</b>	10419
<b>Cytogenetics:</b>	14q11.2
<b>Protein Families:</b>	Stem cell - Pluripotency
<b>MW:</b>	61.1 kDa
<b>Gene Summary:</b>	<p>This gene encodes an enzyme that belongs to the methyltransferase family. The encoded protein catalyzes the transfer of methyl groups to the amino acid arginine, in target proteins that include histones, transcriptional elongation factors and the tumor suppressor p53. This gene plays a role in several cellular processes, including transcriptional regulation, and the assembly of small nuclear ribonucleoproteins. A pseudogene of this gene has been defined on chromosome 4. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Sep 2015]</p> <p>Transcript Variant: This variant (4) contains multiple differences in the 5' coding region, compared to variant 1, including initiation of translation at a downstream in-frame start codon. The encoded isoform (d) has a shorter N-terminus than isoform a.</p>